TYLOSIN-20%

Oral Solution For Poultry, Turkey, Pigeons



COMPOSITION: Each ml contains: Tylosin (As Tylosin Tartrate)

200 mg

PROPERTIES:

- Tylosin as a strong macrolide antibiotic covers a wide spectrum of Gm +ve bacteria such as: Staph, Strept, Diplococcus, Coryne, Clostridia and other bacterial infections secondary to viral diseases.
- Very effective against mycoplasma infections and affects also some riclettsies (Group psittacosischlamydiosis) and Haemophilus gallisepticum.
- Some Gm-ve bacteria (Pasteurella, Campylobacter) also are affected by tylosin.

INDICATIONS:

- For prevention and treatment of: Mycoplasmosis, CRD, Air succulitis, Infectious Synovitis, Infectious Sinusitis, Chlamydiosis, Spirochaetosis.
- Stress conditions due to vaccination, debeaking, transporting, rehousing, changes of feed and weather.

DOSAGE:

- Respiratory or general infections in poultry: 500 ml per 200 litres of water for 3-5 days according to severity of the disease.
- Spirochaetosis: (2.5% solution), used as preventive and for control.
- Pigeons: 0.075-0.125 ml of the product / kg B.wt. / 4 times daily. (15-25 mg activity / kg B.wt.)

PREVENTION PROGRAM:

- Use 500 ml per 200 litres of water as follows:
 - **Broliers:** 1st week of age for 3 days.
 - 4th week of age for 1 day.
 - Pullets: 1st week of age for 3 days. 4th week of age for 1 day. 9th week of age for 2 days. 20th week of age for 2 days.
 - **Parent stocks:** 1st week of age for 5 days.
 - $4^{th},\,9^{th},\,16^{th}$, 20^{th} and 24^{th} week of age for 2 days.
 - **Turkeys:** 1 st week of age for 5 days. 3rd week of age for 1 day.

REMARK: Not recommended for table-eggs layers.

During treatment, eggs not used for human consumption.

SAFETY PERIOD: For meat consumption: 7 days.

STORAGE: Keep in a dry, dark place below 30 °C.

PACKING: (100, 250, 500, 1000) ML.