

TYLOSIN-50%

Water Soluble Powder
For Poultry, Turkeys, Pigeons

VAPCO

ANTIBACTERIAL

COMPOSITION: Each gm contains:

Tylosin (As Tylosin Tartrate)

500 mg

PROPERTIES:

- Tylosin as a strong macrolide antibiotic covers a wide range of Gm+ve bacteria such as: Staph, Strept, Diplococcus, Coryne, Clostridia and other bacterial infections secondary to viral diseases.
- Very effective against mycoplasma infections and affects also some rickettsias (Group psittacosis – chlamydiosis) and Hemophilus gallisepticum.
- Some Gm-ve bacteria (Pasteurella, Campylobacter) also are affected by Tylosin.

INDICATIONS:

- For prevention and treatment of: Mycoplasmosis (CRD), Air sacculitis, Infectious synovitis, Infectious sinusitis, Chlamydiosis, Spirochaetosis.
- Stress conditions due to vaccination, debeaking, transporting, rehousing, changes of feed and weather.

DOSAGE:

- **Solution preparation:** Add 2 litres of water to Tylosin 50% (not vice versa) and dissolve well then add to the water tank.
- **Treatment:**
 - Respiratory or general infections in poultry: (0.05% solution) 200 gm of tylosin 50% per 200 litres of water for 3-5 days according to severity of the disease.
 - Spirochaetosis: (0.5% solution) used as preventive and for control.
 - Pigeons: 0.03 – 0.05 gm/ kg b.wt . 4 times a day
(15-25 mg activity / kg B.wt. / 4 times daily).
- **Prevention program:**
Use 200 gm per 200 litres of water as follows:
 - **Broilers:** 1st week of age for 3 days.
4th week of age for 1 day.
 - **Pullets:** 1st week of age for 3 days.
4th week of age for 1 day.
9th and 20th week of age for 2 days.
 - **Parent stocks:** 1st week of age for 5 days.
4th, 9th, 16th, 20th, and 24th week of age for 2 days.
 - **Turkeys:** 1st week of age for 5 days.
3rd week of age for 1 day.

SAFETY PERIOD: For meat consumption: 7 days.

REMARK:

- Not recommended for table-eggs layers.
- During treatment , eggs not used for human consumption.

STORAGE: Keep in dry, dark place below 30 °C.

PACKING: 113 gm and others.