


# TYLOVAP

Injectable Solution

For Sheep, Goats, Cattle, Pigs, Dogs, Cats , Poultry



**COMPOSITION:** Each ml contains:  
**Tylosin (As Tylosin Tartrate) 200 mg**

## PROPERTIES:

A macrolide antibiotic which is mainly active against Gm+ve, some Gm-ve bacteria, large viruses, certain spirochaetes and mycoplasma.

## INDICATIONS:

For treatment of the followings:

- Respiratory and uro-genital tracts infections, conjunctivitis, tonsillitis, otitis externa, metritis, rhinitis anthrites.
- Mycoplasma, calf diphtheria, vibrio, foot rot, listeria, contagious agalactia, shipping fever, leptospira, erysipelas.
- Enteritis, dysentery, arthritis.
- Infections of Chlamydia, clostridia, staph, strept, etc...
- Wounds, pyodermitis and secondary infections during viral diseases.
- Mastitis (supportive treatment).
- Poultry: Coryza, CRD, sinusitis, spirochaeta, pasteurella, synovitis, air succulitis.

## DOSAGE:

Daily by intramuscular injection for 3-5 days.

- **Sheep, goats, dogs, cats:** 0.5-1.0 ml / 10 kg B. wt.
- **Cattle:** 2-5 ml / 100 kg B.wt..
- **Calves, Pigs:** 0.5-2.5 ml / 50 kg B.wt..
- **Poultry:** By subcutaneous injection: 0.05-0.1 ml / kg B.wt.
- **Turkeys (Sinusitis):** Inject into sinuses (0.12-0.25 ml). Repeat after 10 days when necessary.

## REMARK:

- Inject only (10) ml I.M. at one site.
- Not used for table- eggs layers
- During treatment, eggs not used for human consumption.

## SAFETY PERIOD:

- **Poultry meat & eggs:** 5 days.
- **Milk:** 4 days.
- **Swine meat :** 14 days.
- **Other meat:** 21 days.

**STORAGE:** Keep in dry, dark place below 30 °C..

**PACKING:** (50, 100)ml.