

VACOMIL-35%

**FUNGICIDE
WETTABLE POWDER**

VAPCO

Class:
Acylalanine

A systemic fungicide effective in controlling fungal diseases such as: Blights, Downy mildew and soil-borne diseases.

COMPOSITION: Each kg contains:

METALAXYL

35% W/W (A.I)

PROPERTIES:

VACOMIL-35%: A systemic fungicide effective in controlling fungal diseases on a wide range of crops.

VACOMIL-35%: Used on many crops such as: Potatoes, Cucurbits, Onion, Garlic, Grape, Strawberry, Fruit trees, Olives, Ornamentals and soil-borne diseases.

VACOMIL-35%: Used as seed treatment to protect from soil diseases.

USES AND APPLICATION RATE:

CROP	DISEASE	APPLICATION RATE GM/ 20 LT. WATER	SAFETY PERIOD (DAYS)
Potato, Tomato, Cucumber, Pumpkin, Squash, Watermelon.	Downy Mildew, Late Blight	15	7
Onion, Garlic.	Downy Mildew, Blight, Mold	6 - 12	7
Strawberries	Leaf Spot, Root Rot, Stem Rot	15	15
Grapes	Downy Mildew	15 - 20	7
Apricot, Plum, Cherry, Peach, Almond, Apple, Pear.	Leaf Spot, Leaf Curl, Scab	20	15
Olives	Leaf Spot	8-10	15
Ornamentals	Downy Mildew	8-10	---

CAUTION:

Used only on above mentioned crops and follow the recommended safety period before harvest.

NOTICE:

Used as seed dressing for the control of emergence diseases used at rate 200 gm product / 100 Kg seeds.

SLURRY TREATMENT:

Carefully mix the appropriate quantity of **VACOMIL-35%** for the corresponding quantity of seeds with sufficient water to obtain fluid mixture, for 3-5 minutes.

DRY TREATMENT:

Used a mechanical mixer, place the seed together with 0.5 – 1 lt. water/100 Kg seeds, shake for 1-2 minutes, then add the corresponding dose of **VACOMIL-35%**, and re-shake for about 3 minutes.

• For more details about first aid & precautions please refer to first aid & precautions index.

RE-ENTRY PERIOD:

48 hours after the last application.

COMPATIBILITY:

Compatible with most other pesticide except alkaline compounds.

• For more details about first aid & precautions please refer to first aid & precautions index.