

# VACOMIL-EXTRA 44%

**FUNGICIDE  
SUSPENSION CONCENTRATE**

**VAPCO**

Class:  
**Chloronitrile<sup>(1)</sup>  
Acylalanine<sup>(2)</sup>**

A systemic and protectant fungicide used to control many of fungal diseases on a wide range of agricultural crops.

**COMPOSITION:** Each liter contains:

**CHLOROTHALONIL<sup>(1)</sup>**

**40% W/V (A.I)**

**METALAXYL-M (MEFENOXAM)<sup>(2)</sup>**

**4% W/V (A.I)**

## **PROPERTIES:**

**VACOMIL-EXTRA 44%:** Systemic fungicide with protective action.

**VACOMIL-EXTRA 44%:** Used to control many of fungal diseases on a wide range of agricultural crops.

**VACOMIL-EXTRA 44%:** Rapidly taken by plant tissue and distributed within leaves and stem.

**VACOMIL-EXTRA 44%:** Inhibiting fungal growth within the plant and provides a residual protective film over the surface of plant that inhibiting spore germination.

## **USES AND APPLICATION RATE:**

<b>CROP</b>	<b>DISEASE</b>	<b>APPLICATION RATE ML / 100 LT. WATER</b>	<b>SAFETY PERIOD (DAYS)</b>
Tomato, Potato.	Early Blight, Late Blight.	250	14
Watermelon, Muskmelon, Cucumber, Squash, Pumpkin.	Downy Mildew, Leaf Spot, Gummy Stem Blight, Alternaria Leaf Spot.	200	7
Cauliflower, Cabbage, Broccoli.	Downy Mildew, Alternaria Leaf Spot, Ringspot.	200	14
Brussels Sprout.	Downy Mildew, Alternaria Leaf Spot, Ringspot.	200	14
Broad Beans.	Downy Mildew.	200	14
Onions, Garlic, Leeks.	Downy mildew, Botrytis, Purple Blotch.	200	14
Raspberry, Blackberry, Boysenberry.	Botrytis, Dryberry, Cane spot.	375	14
Grapes	Black spot, Botrytis, Downy mildew, Powdery Mildew.	375	28
Ornamentals	Downy mildew, Botrytis, White Rust.	200	---

## **CAUTION:**

Used only on above mentioned crops and follow the recommended safety period before harvest.

## **RE-ENTRY PERIOD:**

48 hours after the last application .

## **COMPATIBILITY:**

Compatible with most pesticides, except highly alkaline materials.

• For more details about first aid & precautions please refer to first aid & precautions index.