

VACOMIL-MZ 72%

FUNGICIDE
WETTABLE POWDER

VAPCO

Class:
Dithiocarbamate⁽¹⁾
Acylalanine⁽²⁾

Effective fungicide acts as systemic and contact fungicide to control a wide range of fungal diseases such as Blights and Downy mildew on a wide varieties of crops.

COMPOSITION: Each kg contains:

MANCOZEB⁽¹⁾

64% W/W (A.I)

METALAXYL⁽²⁾

8 % W/W (A.I)

PROPERTIES:

VACOMIL-MZ 72%: A special formulation of Mancozeb and Metalaxyl which protect plant in development growth stage, Metalaxyl absorbed through the tissue of plant, while Mancozeb protect outside the tissue.

VACOMIL-MZ 72%: Control a wide range of fungal diseases on a wide varieties of crops.

VACOMIL-MZ 72%: Has an immediate effect against diseases such as Blights, Downy mildew diseases on Grapes, Tomatoes, Potatoes, Onion and other crops which mentioned in below table, early and late blight on tomato, potato and have affected in leaf spot diseases.

USES AND APPLICATION RATE:

CROP	APPLICATION RATE GM/ 20 LT. WATER	TIME BETWEEN APPLICATION	NOTE	SAFETY PERIOD (DAYS)
Tomatoes (on field)	70	7 days	3 Times during growing period first application when first symptoms appear during active growing period. When plant grow and mature use Another common fungicide during 7 days from last VACOMIL-MZ 72% spraying.	7
Potatoes	50	7 days	3 Times during active growing period, another fungicide could be used after 7 days from Last VACOMIL-MZ 72% application.	14
Onion	50	7 days	3 Times maximum, first application before first symptoms of disease appears.	10
Melon, Squash	50	14 days	---	4
Lettuce	50	14 days	---	7
Pepper	50 – 60	21 days	---	7

• For more details about first aid & precautions please refer to first aid & precautions index.

CROP	APPLICATION RATE GM/ 20 LT. WATER	TIME BETWEEN APPLICATION	NOTE	SAFETY PERIOD (DAYS)
Grapes	50	10 days	Four Times maximum of applications: 1. When the plant height is 20-25 cm. 2. Before flowering stage after 14 days of the first application. 3. At end of flowering period. 4. After 14 days of complete flowering stage, it is preferable to use other common fungicide through 7 days from last VACOMIL-MZ 72% application.	14
Fruit trees	140	First application in spring and the second one in fall after harvest	Width of the furrow is about 90 Cm ²	7
Citrus trees	60 – 120	3 months	---	21

CAUTION:

Used only on above mentioned crops and follow the recommended safety period before harvest.

RE-ENTRY PERIOD:

48 hours after the last application.

COMPATIBILITY:

Compatible with most known fungicides and insecticides.

*** OTHER FORMULATION & TRADE NAMES: VACOMIL-FORTE 58% WP (MANCOZEB 48% + METALAXYL 10%), PANALAXIL – 72% WP.**

• For more details about first aid & precautions please refer to first aid & precautions index.