

VACOMIL-PLUS 50%

**FUNGICIDE
WETTABLE POWDER**

VAPCO

Class:
Acylalanine⁽¹⁾
Inorganic compound⁽²⁾

Effective systemic preventive and curative fungicide to control a wide range of fungal diseases such as (Blight, Downy mildew) on a wide range of crops and medical plants.

COMPOSITION: Each kg contains:

METALAXYL⁽¹⁾

15% W/W (A.I)

COPPER OXYCHLORIDE⁽²⁾

35% W/W (A.I)

PROPERTIES:

VACOMIL-PLUS 50%: Is a curative and preventive fungicide to control a wide range of fungal diseases on a wide range of crops.

VACOMIL-PLUS 50%: Is effective against a wide range of fungal diseases that affected a wide range of crops.

VACOMIL-PLUS 50%: Not phytotoxic for treated plants except some species of Apples.

VACOMIL-PLUS 50%: Controls a wide range of diseases such as: Downy mildew, Blights, Leaf spot, Scab.

VACOMIL-PLUS 50%: Is also used as seed treatment for number of crops.

USES AND APPLICATION RATE:

CROP	DISEASE	APPLICATION RATE GM/ 20 LT. WATER	SAFETY PERIOD (DAYS)
Cauliflower, Cabbage	Downy mildew.	30 – 35	14
Watermelon, Potato, Muskmelon, Tomato	Early Blight, Late Blight.	30 – 35	14
Apple, Pear	Downy Mildew, Scab, Canker	30 – 35	21
Peach, Apricot, Plum, Cherry	Downy Mildew, Scab, Canker	30 – 35	21
Grapes	Downy Mildew.	30 – 40	12
Medicinal plant	Late Blight, Downy Mildew.	30	20

CAUTION:

Used only on above mentioned crops and follow the recommended safety period before harvest, some varieties of red Apple are sensitive to this product. Sensitivity test should be done before application.

• For more details about first aid & precautions please refer to first aid & precautions index.

RE-ENTRY PERIOD:

48 hours after the last application.

COMPATIBILITY:

Compatible with most other pesticides, but incompatible with Mercuric compounds, DNOC, Lime sulfur and Dithiocarbamates.

*** OTHER FORMULATION:**

VACOMIL-PLUS – 76.5% WP

(METALAXYL 15% + COPPER OXYCHLORIDE 61.5% (Cu 35%)).

• For more details about first aid & precautions please refer to first aid & precautions index.