

VAPCOMIC-1.8%

**INSECTICIDE / ACARICIDE
EMULSIFIABLE CONCENTRATE**

VAPCO

**Class:
Avermectin**

Effective insecticide and acaricide by contact and stomach action.

COMPOSITION: Each liter contains:

ABAMECTIN

1.8% W/V (A.I)

PROPERTIES:

VAPCOMIC-1.8%: Can provide more effect, longer control of all varieties of Spider mites, Cyclamen mite, Rust mite, Leafminer, Psylla, Thrips, Worm..... etc at all stages on a wide range of crops.

VAPCOMIC-1.8%: Highly active for a long period due to its content of a high percent of the active isomer avermectin B1a.

USES AND APPLICATION RATE:

CROP	PEST	APPLICATION RATE ML / 20 LT. WATER	SAFETY PERIOD (DAYS)
Tomatoes	Leafminers, Pinworm, Red Spider mites Rust Mites.	5 - 7	3 - 7
Cucumber, Eggplant	Leafminers, Red Spider mites.	5 - 7	3 10
Cucurbits	Leafminers, Thrips, Mites	20 - 40	3
Pepper, Celery, Melon, watermelon	Leafminers, Mites	10 - 20	7
Lettuce	Leafminers, Mites	20 - 40	14
Cabbage	Diamondback Moth	40	7
Potato	Leafminers, Thrips, Pyslla, Worms	20	7
Radish	Mites	25	14
Onions, Garlic	Thrips, Leafminers, Mites.	5 - 10	3
Squash	Leafminers	16 - 32	3
Chicory	Mites	16	7 - 14
Endive	Mites	16	28

• For more details about first aid & precautions please refer to first aid & precautions index.

CROP	PEST	APPLICATION RATE ML / 20 LT. WATER	SAFETY PERIOD (DAYS)
Strawberries	Two Spotted Spider mite, Cyclamen Mite, Red Spider mites.	5 - 7	7
Citrus	Leafminers, Red Spider mites.	10	7
Almond	Mites, Pyslla	25	21
Apple, Walnut	Mites	25 - 48	21
Pears	Mites	25 - 48	10
Date Palm	Mites	10	10
Avocado	Mites	15	14
Banana	Thrips	90	---
Papaya	Mites	6 - 24	14
Pineapple	Mites	10 - 40	---
Cotton	Mites	10 - 20	7
Rice	Mites	2 - 3 lt/ha	3
Ornamentals	Leafminers, Red Spider mites.	5 - 7	3 - 7

CAUTION:

Used only on above mentioned crops and follow the recommended safety period before harvest.

REMARKS:

- * Do not spray the crop more than 5 times during one season.
- * Results of control appears after 3-5 days of spraying, however effect of the product appears immediately after spraying.
- * Quantity of the solution must not exceed 110 ml of **VAPCOMIC** /1000 m² for vegetables and ornamentals.

RE-ENTRY PERIOD:

Do not enter field before 24 hours from last application under field conditions, the spray should be allowed to dry on the foliage before re-entry into the treated areas.

PHYTOTOXICITY:

Non-phytotoxic, but do not spray it on carnations and sedges, toxic to Pome fruits when mixed with Captan (fungicide) containing compounds.

COMPATIBILITY:

Compatible with most pesticides, but compatibility test is recommended before mixing.

*** OTHER FORMULATION AND TRADE NAMES: ABAMEX-1.8% (EC, SL), BAMECTIN-1.8% (EC, SL), MAYSAMECTIN-1.8% EC, VAPCOMIC-2% EC W/W.**

• For more details about first aid & precautions please refer to first aid & precautions index.