# **VAPCOMIC GOLD-8.4%**

# ACARICIDE/INSECTICIDE SUSPENSION CONCENTRATE



Is effective acaricide and insecticide by contact and stomach action used to control a wide range of insects in motile stages especially adult stage which is difficult to control.

**COMPOSITION:** Each liter contains:

**ABAMECTIN** 

8.4% W/V (A.I)

**PROPERTIES:** 

**VAPCOMIC-GOLD:** Is effective acaricide and insecticide by contact and stomach action used to control a wide range of insects in the motile stages especially adult stage which is difficult to control.

**VAPCOMIC-GOLD:** Highly active for a long period due to its content of a high percent of the active isomer Avermectin B1a.

### **USES AND APPLICATION RATE:**

CROP	PESTS	APPLICATION RATE ML / 20 LT. WATER	SAFETY PERIOD (DAYS)
Tomatoes, Chili, Eggplant. Potato.	Leafminer larvae, Red spider mites Rust mites, Thrips, Tuber moth.	3	3 14
Cucumber, Squash Melon & Water Melon	Red spider mites, Thrips.	3	7
Pepper, Celery	Leafminers, Mites.	2.5 - 4.5	7
Lettuce	Leafminers, Mites.	4.5 - 8.5	14
Cabbage	Diamondback Moth	8.5	7
Radish	Mites	5	14
Onions, Garllic	Thrips, Leafminers, Mites.	3	3
Dry Beans	Leafminers, Red Spider mites.	3 - 5	7
Chicory	Mites	3.5	7 - 14
Endive	Mites	3.5	28
Strawberries	Two-spotted Spider mite, Cyclamen Mite, Red Spider mites.	3	3

<sup>•</sup> For more details about first aid & precautions please refer to first aid & precautions index.

CROP	PESTS	APPLICATION RATE ML / 20 LT. WATER	SAFETY PERIOD (DAYS)
Citrus	Leafminers, Red Spider mites, Rust Spider mite, Thrips.	2.5	10
Grapes	Pacifi C Spider Mite, Two-spotted Spider Mite, Variegated Leafhopper, Western Grape Leafhopper, Western Grapeleaf Skeletonizer,Willamette Spider Mite.	3 - 5	28
Almond	Mites, Pyslla	5	21
Apple, Walnut	Mites	5 - 10	21
Pears	Mites	5 - 10	10
Date Palm	Mites	3	10
Avocado	Mites	3	14
Banana	Thrips	19	
Papaya	Mites	3 - 5	14
Pineapple	Mites	3 - 8	
Cotton	Mites	3 - 4	7
Rice	Mites	430-640 MI/Ha	3
Hops	Two-spotted Spider mite	3 - 5	28
Mint	Two-spotted Spider mite	2.6 - 3.7	28
Ornamentals	Leafminers, Red Spider mites.	2.5 3	

Hectare= 10000 m<sup>2</sup>

### CAUTION:

Used only on above mentioned crops and follow the recommended safety period before harvest.

# **REMARKS:**

- Do not spray the crop more than 5 times during one season.
- Results of control appears after 3 5 days of spraying, however effect of the product appear immediately after spraying.
- Quantity of the solution must not exceed 23 ml of VAPCOMIC GOLD / 1000 m<sup>2</sup> for vegetables and ornamentals.
- For mites & leafminers control: spray both surfaces of the leaf once the injury appears
- When spraying citrus tree: avoid overspray when using VAPCOMIC GOLD with paraffin oil.

## **RE-ENTRY PERIOD:**

Do not enter field before 48 hours from last application, the spray should be allowed to dry on the foliage before re-entry into treated areas.

# PHYTOTOXICITY:

Not-phytotoxic, at the recommended rates but do not spray it on carnations and sedges, and phytotoxic to Pome fruits when mixed with Captan (fungicide) compounds.

#### COMPATIBILITY:

Compatible with most pesticides, except captan compound but compatibility test is recommended before mixing

• For more details about first aid & precautions please refer to first aid & precautions index.