


VAPCOTRIM-PLUS Potentiated

**Water Dispersible Powder
For Poultry & Livestock**



COMPOSITION: Each gm contains:

Erythromycin Thiocyanate	180 mg
Trimethoprim	30 mg
Sulphadiazine - Sodium	150 mg

PROPERTIES:

- Erythromycin inhibits the protein synthesis of bacteria, while trimethoprim magnifies the effect of sulphonamide by blocking transformation of folic acid into bio-active folic acid.
- Effective against Gm+ve and Gm-ve bacteria: Coli, Staph, Strept, Salmonella, Pasteurella, Vibrio, Mycoplasma, Clostridia, Listeria, Coryne etc...

INDICATIONS:

- **Poultry:**
 - For prevention and treatment of: CRD, Mycoplasmosis, Coryza, Infectious sinusitis, Air sacculitis, Infectious synovitis, Vibronic hepatitis, Salmonellosis, Pasteurellosis, Colibacillosis, enteritis, diarrhea.
 - Stress conditions due to vaccination, debeaking, rehousing, transporting, changes of feed and weather.
- **Other animals:** Cases of pneumonia, bronchitis, enteritis and diarrhea etc..

DOSAGE:

Mix well before use.

- **Treatment:** (Duration for 5 days).
 - **Poultry:** 100 gm per 200 litres of drinking water (use hot water 50-60 °C).
 - **Livestock:** 2 gm per 15 kg B.wt. daily and orally with water.
- **Prevention:** Half dosage for the same period.

REMARKS:

- Flumequine should not be associated with trimethoprim-sulphonamide treatment.
- Not recommended for laying flocks.
- Contra-indicated in cases of liver and Kidney troubles, Gumboro disease.
- Adequate water intake must be available to avoid Sulphonamide precipitation.

SAFETY PERIOD: For meat and milk consumption: 7 days.

STORAGE: Keep in a dry, dark place below 30 °C.

PACKING: (100, 250, 500, 1000) GM.