


VAPCOVIT

Water Dispersible Powder
Vitamins For Poultry And Livestock



COMPOSITION: Each gm contains:

| | | |
|-------------------------|--------|------|
| Vitamin A | 20,000 | I.U. |
| Vitamin D3 | 5000 | I.U. |
| Vitamin E | 5.0 | I.U. |
| Vitamin K3 | 3 | mg |
| Vitamin C | 20 | mg |
| Vitamin B1 | 1.5 | mg |
| Vitamin B2 | 2 | mg |
| Vitamin B6 | 2 | mg |
| Vitamin B12 | 0.02 | mg |
| Nicotinic acid | 20 | mg |
| Calcium-D- Pantothenate | 6.5 | mg |
| Folic acid | 0.4 | mg |

INDICATIONS:

- Hypovitaminosis.
- Stress conditions due to vaccination, debeaking, transporting, rehousing, changes of feed and weather.
- Maintains good performance and increases the production.
- Improvement of feed conversion, productivity, the growth, fertility and hatchability rates, good feathering and flesh development.
- Enhancement of better rearing results.
- Helps in producing high eggs-quality.
- Increase of viability and resistance against diseases.
- Effective in cases of caged-layers fatigue.

DOSAGE: Duration for (3-5) days.

Mix well before use.

- **Poultry:** 100 gm per 200 litres of drinking water.
- **Calves, Foals, (Weaners):** 2 gm per animal daily and orally with water.
- **Calves, Foals (Suckling):** 1 gm per animal daily and orally with water.
- **Via Feed:** 1 kg per ton of feed.

STORAGE: Keep in a dry, dark place below 25 °C.

PACKING: (100, 250, 500, 1000) GM.