

# VAPCOZIN –20 EC

Insecticide And Acaricide  
Emulsifiable Concentrate

**VAPCO**  
INSECTICIDES

**COMPOSITION: Each Liter contains:**

**Amitraz**

**200 gm**

## PROPERTIES:

- Non-systemic insecticide and acaricide of the amidine group with contact and respiratory action.
- Characterized by its strong effect against ticks which are resistant to arsenical, organo-phosphorus and organo-chlorine insecticides.
- Rapidly degraded in the soil and does not cause environmental pollution hazards.
- Low mammalian toxicity.
- Expellent action causes ticks to withdraw mouth parts rapidly and fall off the host animal.

## USES:

- Treatment of different strains of scab and mange.
- Control of many species of ticks, mites, lice, keds, fleas and flies on sheep, goats, cattle, camels and dogs.
- For combating insects in farms, stables and other animals houses.
- Destruction of flies-larvae in breeding sites and keeping down flies population.

## METHOD OF APPLICATIONS:

By spraying or dipping after dilution with water as follows:

Species	Initial dilution rate	Interval	Replenishment
<b>Cattle, Camels:</b>			Replenishment rate is half of the initial dilution rate
Ticks, Lice,	1:1100	7-10 days	
Mange	1:800	12 days	
<b>Sheep, Goats</b>			
Ticks,	1:1100	At need	
Scab	1:400	At need	
Keds, Lice	1:400	10-12 days	
<b>Dogs:</b>			
Ticks, Mange	1:400	10 days	
<b>Animals Houses</b>	1-2 %	At need	

## PRECAUTIONS:

- Toxic to equines and fishes.
- Avoid contact with eyes, mouth, skin and inhalation.
- Before spraying, wear protective clothes, gloves, boots and goggles.
- Do not eat, drink or smoke during spraying.
- Avoid polluting ponds, water ways and water sources.
- Do not allow off-spring to suckle until mothers are completely dried from the insecticide liquid.



**VAPCO**  
**INSECTICIDES**

**FIRST – AID:**

- Eye contact: wash with water for 15 minutes.
- Skin contact: wash with soap and water.
- Inhalation:
  - Move patient to fresh air
  - Apply artificial respiration, give oxygen when necessary.
- Ingestion:
  - Do not induce vomiting.
  - wash the mouth with water.
  - let patient, if conscious , to drink 1-2 glasses of water to dilute the product in the stomach.
  - Apply gastric lavage.

Get medical help, when necessary

**ANTIDOTE:** Not specific, treat according to symptoms.

**SAFETY PERIOD:**

- For milk consumption: 2 days.
- For meat consumption: Sheep ,Goats: 7 days.  
Cattle: 1 day.

**STORAGE:**

- Keep away from children, food and drinks.
- Keep in a dry, dark place below 30 °C.

**PACKING:** 250 ml

(1, 5, 20) Litres.