

VAYOMIL - 24%

**INSECTICIDE, ACARICIDE AND NEMATOCIDE
SOLUBLE CONCENTRATE**

VAPCO

**Class:
Carbamate**

Contact and systemic insecticide, acaricide, and nematocide. to control wide range of pests on many crops .

COMPOSITION: Each Liter contains:

OXAMYL

24% W/V (A.I)

PROPERTIES:

VAYOMIL-24%: Contact and systemic insecticide, acaricide, and nematocide. absorbed by the foliage and roots, with translocation.

VAYOMIL-24%: Completely systemic pesticide translocation occurs within leaves to the roots to protect the plant from mites, insects and nematode.

VAYOMIL-24%: Highly toxic to mammals, toxic to fish and bees.

USES AND APPLICATION RATES:

CROPS	PESTS	APPLICATION RATE ML/20 LT WATER	SAFETY PERIOD (DAYS)	NOTES
Tomatoes, Carrots, Cucumbers, Melons, Watermelons, Pumpkin, Potatoes, Green Pepper.	Canker Nematodes, Beetles, White Flies, Thrips, Leafminers & Mites.	50 (SPRAY) 50 (DRENCH)	3	To control nematodes spray the vegetative growth after 2 weeks from transplanting, re-spray after 3 weeks.
		50	8	
Field Crops, Cereals And Tobacco.	Root-Knot Nematodes, Canker Nematodes, Cyst Nematode, Stunt Nematode, Aphids, Thrips, Leafminers And Whiteflies	50	21	Re-spray after 3 weeks if necessary.
Apples, Pears, Orange, Lemon, Mandarin, Pummelo , Grapefruit.	Root-Knot Nematodes, Canker Nematodes, Whiteflies, Thrips, Leafminers, Aphids And Red Spidermites.	50 (SPRAY) 50 (DRENCH)	14	

• For more details about first aid & precautions please refer to first aid & precautions index.

CAUTION:

Used only on above mentioned crops and follow the recommended safety period before harvest.

REMARKS:

- Use **VAYOMIL-24% SL** by irrigation systems, or spraying on the soil or by drop irrigation system or foliar spraying.
- Add phosphoric acid to reduce pH and get best result.

RE-ENTRY PERIOD:

72 hours after the last application.

COMPATIBILITY:

Compatible with most common pesticides except alkaline compounds.

• For more details about first aid & precautions please refer to first aid & precautions index.