


VAPCOPHORE - 0.5%

Solution
Disinfectant And Sanitizer



COMPOSITION :

An iodine polyaldehyde carbonic acid complex with ortho-phosphoric acid in aqueous solution containing 2.5 % iodophore (Equivalent to 0.5 % active iodine).

PROPERTIES:

- Has a broad spectrum effect against covered and naked viruses especially adeno-viruses.
- Highly effective against most bacteria, protozoa, yeast, fungi and algae.
- Harmless to skin and soft tissues.

USAGE:

After dilution with water as follows:

- 2-4% For cleaning and disinfection of stables, equipment, feeding troughs, pens, milking machines, hatcheries and processing plants.
- 1-2 % For poultry farms and hatching eggs dipping.
- 8-12 % For control of epidemics (Salmonellosis, Brucellosis, Newcastle etc.), and in case of general danger of epidemics.
- 2 % For teat dipping or udder washing as an aid in the control of mastitis and teat sores. (Apply immediately after milking and before the next milking).
- 0.04% For sanitization of the drinking water of poultry and large animals.

FIRST AID:


- **Ingestion:**
 - Wash the mouth with water.
 - Drink two glasses of water.
 - Do not induce vomiting.
- **Eye contact:** Flush with water for at least 15 minutes.
- **Skin contact:** Wash with water.
- **Inhalation:** Remove to fresh air; give oxygen when necessary.
- **Get medical attention when necessary.**

REMARKS:

- Harmful to fish.
- Don't mix with other chemicals.

CAUTIONS:

- Avoid contact with eyes, skin, inhalation and swallowing.
- Wash containers with water before discarding.



STORAGE: Keep in a dry, dark place at below 30 °C
Keep away from children, food and drinks.

PACKING: (100,250, 500, 1000) ml & 5 litres gallons.