

LEVOZAN BOLUS

Anthelmintic Bolus For Ruminants



COMPOSITION: Each bolus contains:

| | |
|-----------------------|----------------|
| Levamisole HCL | 1000 mg |
| Oxyclozanide | 1400 mg |

PROPERTIES:

- Oxyclozanide removes practically all flukes (*Fasciola* spp.) present in the bile ducts.
- Levamisole is the racemic mixture of optical isomers of tetramisole .
 - * Levamisole removes mature and developing immature stages of wide range of nematodes, and is highly effective against the following species:
 - **In the abomasum:** *Haemonchus*, *Ostertagia* and *Trichostrongylus* spp.
 - **In the intestines:** *Trichostrongylus*, *Cooperia*, *Nematodirus*, *Oesophagostomum*, *Chabertia* and *Bunostomum*.
 - **In the lungs:** *Dictyocaulus* spp., including inhibited *D.viviparus* larvae.

INDICATIONS:

Levozan bolus is a broad spectrum anthelmintic for the treatment and control of gastro-intestinal and pulmonary nematode infestations ,liver chronic fascioliasis ,dicrocoelium (Lancet Flukes, and rumen flukes (*Paramphistomum*) .

DOSAGE:

- Levozan bolus is administered orally, by means of balling gun, or dispersed in water and given orally.
- **Cattle, Camel :** One bolus per 150 kg B.wt.
For animals weighing 300 kg or more, a maximum of 2 boluses should be given.
As the bolus is scored, it may be broken in half to enable the dose to be administered in units of 75 kg B .wt.
- **Sheep, goats:** Half bolus per 50 kg B.wt

SAFETY PERIOD:

- For meat consumption: 14 days after treatment.
- For milk consumption: 24 hours after treatment.

STORAGE: Keep in a dry, dark place below 30 °C

PACKING: 50 Boluses.