# **CONAZOLE 25%**





Is a broad spectrum, systemic fungicide with preventive and strong curative action used to control leaf spot diseases, powdery mildews, rusts and scab on crops mentioned in the table of uses.

**COMPOSITION:** Each liter contains:

DIFENOCONAZOLE 25% W/V (A.I)

PROPERTIES:

CONAZOLE-25%: A broad spectrum, systemic fungicide used to control fungal diseases

including leaf spot, powdery mildew, rusts and scab, affecting wide range of

crops mentioned in the table of uses.

CONAZOLE-25%: Has outstanding curative efficacy and long-lasting preventive disease

control means fewer applications.

CONAZOLE-25%: Is safe to natural enemies and used for IPM programs.

#### USES AND APPLICATION RATE:

| CROP  | DISEASE   | APPLICATION RATE<br>ML/ 20 LT. WATER | SAFETY PERIOD<br>(DAYS) |
|---|---|--------------------------------------|-------------------------|
| Citrus  | Black Rot of Navel Orange                             | 10                                   | 21                      |
| Apples, Apricots,<br>Plums, Peaches,<br>Nectarine, Pears<br>and Almonds | Powdery mildew, scab, rust and leaf spot.             | 8 - 10                               | 14                      |
| Grapes  | Powdery mildew, Gray mold and Black rot               | 8 - 12                               | 14                      |
| Mango   | Anthracnose   | 10                                   | 15                      |
| Tomato  | Early blight, Leaf mold, Powdery mildew and Leaf spot | 10 - 13                              | 7                       |
| Potato  | Early blight  | 10                                   | 14                      |
| Cucurbits   | Gummy stem blight, Powdery mildew and Anthracnose     | 8 - 10                               | 3                       |
| Pepper  | Leaf Spot and Powdery mildew                          | 8 - 12                               | 3                       |
| Carrot  | Fire blight   | 8 - 12                               | 14                      |
| Strawberry  | Leaf spots  | 8 - 10                               | 14                      |
| Sugar Beet  | Leaf spots  | 10                                   | 14                      |
| Beans   | Leaf spots, Rust and<br>Anthracnose                   | 8 - 12                               | 14                      |

<sup>•</sup> For more details about first aid & precautions please refer to first aid & precautions index.

| CROP             | DISEASE  | APPLICATION RATE<br>ML/ 20 LT. WATER | SAFETY PERIOD<br>(DAYS) |
|------------------|--|--------------------------------------|-------------------------|
| Peas             | Ascochyta blight                               | 10                                   | 14                      |
| Peanuts          | Leaf spots                                     | 10                                   | 20                      |
| Garlic and Onion | Purple blotch, Rust, Blight and Powdery mildew | 10 - 13                              | 7                       |
| Ornamentals      | Powdery mildew, Scab. Rust<br>And Leaf spot    | 8 - 10                               |                         |

# CAUTION:

Used only on above mentioned crops and follow the recommended safety period before harvest.

## **RE-ENTRY PERIOD:**

24 hours from last application.

## COMPATIBILITY:

Can be mixed with most insecticides and fungicides. but compatibility test is recommended before spray.

<sup>•</sup> For more details about first aid & precautions please refer to first aid & precautions index.