



DIACARB-15% SC
INSECTICIDE
SUSPENSION CONCENTRATE

IS AN INSECTICIDE WITH CONTACT AND INGESTION ACTIVITY. AFFECTED INSECTS CEASE FEEDING , WITH POOR COORDINATION, PARALYSIS AND ULTIMATELY DEATH. IT IS USED FOR BROAD-SPECTRUM CONTROL OF LEPIDOPTERA IN COTTON, VEGETABLES AND FRUIT.

COMPOSITION:

EACH 1 LITER OF DIACARB-15% SC CONTAINS:

INDOXACARB **15% W/V (A.I)**

COMMERCIAL PROPERTIES:

DIACARB - 15% SC: INDOXACARB, THE ACTIVE INGREDIENT BELONGS TO A NEW CLASS OF INSECTICIDE CHEMISTRY KNOWN AS OXADIAZINES. A SPECIFIC AND UNIQUE MODE OF ACTION.

DIACARB - 15% SC: IT HAS A CONTACT AND STOMACH ACTION ON ALL LARVAL STAGES OF LEPIDOPTEROUS PESTS. IT WORKS BY INHIBITING THE FLOW OF SODIUM INTO NERVE CELLS OF THE TARGETED PESTS.

DIACARB - 15% SC: IT PROVIDES YOU WITH A HIGH LEVEL OF PLANT PROTECTION AND INSECT CONTROL.

DIACARB - 15% SC: YET IT'S APPLIED AT VERY LOW RATES, REDUCING THE CHEMICAL LOAD ON THE ENVIRONMENT. LOW TOXICITY TO USERS, CONSUMERS AND MAMMALS IN GENERAL.

DIACARB - 15% SC: HAS SHORT PRE-HARVEST INTERVALS.

USES AND APPLICATION RATE

CROPS	PESTS	APPLICATION RATE ML/20 LT.	SAFETY PERIOD DAYS
CRUCIFERAE	CABBAGE WORM ,	4.0	1 DAY
	CABBAGE BUTTER FLY, LEAF WORM	5.0	
TOMATO , EGGPLANT,PEPPER , CUCUMBER , WATERMELON, SQUASH.	TOMATO FRUITWORM, LEAFMINER	5.0	3DAYS

ISO 9001 & 14001 REGISTERED



VAPCO

VETERINARY & AGRICULTURAL PRODUCTS MANUFACTURING CO. LTD.



GRAPE	GRAPE BERRY MOTH, LEAF HOPPERS.	5.0	14 DAYS
PACH , APPLE , PEAR , APRICOT	APPLE CODLING MOTH	5.0	20 DAYS
POTATO	POTATO TUBER MOTH	5.0	3 DAYS

CAUTION:

USED ONLY ON ABOVE MENTIONED CROPS AND FOLLOW THE RECOMMENDED SAFETY PERIOD BEFORE HARVEST.

RE-ENTRY PERIOD: DON'T ENTER FIELD BEFORE 48 HOURS FROM LAST APPLICATION.