

PENDILIN- 40%

HERBICIDE
SUSPENSION CONCENTRATE

VAPCO

Class:
Dinitroaniline

Selective herbicide used to control annual grasses and broad-leaved weeds in a wide range of crops.

COMPOSITION: Each liter contains:

PENDIMETHALIN

40 % W/V (A.I)

PROPERTIES:

PENDILIN- 40% : Selective herbicide absorbed by leaves and roots.

PENDILIN- 40% : Used against of most annual grasses and broad-leaved weeds.

PENDILIN- 40% : Applied as pre-plant incorporated, pre-emergence, pre-transplanting, or early post-emergence in a wide range of crops which are mentioned in the table.

USES AND APPLICATION RATE :

CROPS	TIME OF APPLICATION	APPLICATION RATE (LT./HA)	COMMENTS
Winter Barley, Winter Wheat	Pre-emergence of the crop to before leaf sheath erect stage	3.3	Do not apply to crops after 30th November.
Spring Barley	Pre-crop emergence	3.3	- Apply after drilling and before emergence. - Do not apply PENDILIN- 40% alone after the end of march unless rainfall is imminent.
Onion, Leek	Pre-crop emergence drilled and transplanted	3.3	- Apply after drilling but before crop or weed emergence. - Transplanted crops (apply pre-transplanting) - Do not apply to any onion or leek crop when heavy rain is forecast.
Carrot , Parsnip	Pre-crop emergence	3.3	- Apply after drilling but before crop and weed emergence.
Broccoli, Cabbage, Cauliflower	Before transplanting	3.3	- Apply after final plant bed cultivation but before transplanting - Do not apply PENDILIN- 40% post-planting as crop damage may occur. - Do not apply PENDILIN- 40% to any transplanted Brassicas when heavy rain is forecast.

• For more details about first aid & precautions please refer to first aid & precautions index.

CROPS	TIME OF APPLICATION	APPLICATION RATE (LT./HA)	COMMENTS
Potato	Pre-emergence of the crop	3.3	- Apply as soon as possible after planting and final ridging up. - Do not apply later than 7 days before emergence.
Strawberry	After flower initiation but before flower trusses emergence	3.3	- PENDILIN- 40% should be applied to dormant newly planted runners in the autumn or early spring immediately after planting and prior to weed emergence. - Pre-planting application of PENDILIN- 40% is not recommended.
Apple, Cherry, Pear, Plum	Before bud burst	3.3	- Apply PENDILIN- 40% to weed-free soil from autumn to early spring. - PENDILIN- 40% is not recommended for use after the end of march unless adequate soil moisture is present

CAUTION:

Hectare = 10000m²

Used only on the above mentioned crops.

NOTES:

- Do not apply **PENDILIN- 40%** to crops suffering from stress.
- Seed should be covered with a minimum of 3.2 cm of settled soil.
- Do not disturb the soil after **PENDILIN- 40%** has been applied as this will result in reduced weed control.
- Do not spray undersown crops.
- **PENDILIN- 40%** should not be used on protected crops, or in greenhouses.
- Do not use on soils with more than 10% organic matter.
- Some soil moisture must be present for **PENDILIN- 40%** to be activated. Best results will be obtained if rainfall occurs within seven days of application.
- All dosages should be applied with 100-200 liters of water per hectare.

RE-ENTRY PERIOD:

Don't enter field before 24 hours from last application.

GRAZING RESTRICTIONS:

Do not graze or feed animal from treated area for 24 hours from last application.

COMPATIBILITY:

Incompatible with most common pesticides and fertilizers.

• For more details about first aid & precautions please refer to first aid & precautions index.