

PROPORID 72.2%

FUNGICIDE
SOLUBLE CONCENTRATE

VAPCO

Class:
Carbamate

Systemic fungicide with preventive and curative action used to control soil born fungi and fungal diseases on crops mentioned in the table of uses.

COMPOSITION: Each liter contains:

PROPAMOCARB HYDROCHLORIDE

72.2% W/V (A.I)

PROPERTIES:

PROPORID-72.2%: Systemic fungicide with preventive and curative protective action used to control soil born fungi and fungal diseases on crops mentioned in the table of uses.

PROPORID-72.2%: Has long residual activity for 3 - 8 weeks depends on temperature, lights and soil type.

USES AND APPLICATION RATE:

CROP	DISEASE	APPLICATION RATE	SAFETY PERIOD (DAYS)
Tomato, Cabbage, Raddish and Potato	Root rot (<i>Pythium</i> and <i>Phytophthora</i>)	Seed planting: Directly after planting, add 70 ml/20 L water to the soil surface at the rate of 2 - 4 L (solution)/m ² .	7 days except cabbage 14 days
	Downy mildew	Nursery: • Before transplanting: add 70 ml/20 L water to the soil surface at the rate of 2 - 4 L (Solution)/m ² . • After transplanting: add 20ml / 20 L water to the soil surface at the rate of 200 ml (Solution) / plant, repeating every two weeks as necessary. Foliar spray: 30 ml / 20 L water	
Cucumber	Late blight (<i>Phytophthora</i>)		
	Root rot (<i>Pythium</i> and <i>Phytophthora</i>)	Seed planting: Directly after planting, add 70 ml /20 L water to the soil surface at the rate of 2 - 4 L (Solution)/m ² .	3 days
	Downy mildew	Nursery: • before transplanting: Add 70 ml /20 L water to the soil surface at the rate of 2 - 4 L (Solution)/m ² . • After transplanting: Add 20 ml / 20 L water to the soil surface at the rate of 200 ml (Solution) / plant, repeating every two weeks as necessary. Foliar spray: 50 ml / 20 L water	

• For more details about first aid & precautions please refer to first aid & precautions index.

CROP	DISEASE	APPLICATION RATE	SAFETY PERIOD (DAYS)
Ornamental	Root rot (<i>Pythium and Phytophthora</i>)	Seed Planting and Nursery: Uses and application rate same as mentioned above. Cutting: Add 5 - 15ml /10 L water to the soil surface at the rate of 5 - 10 L (Solution)/M ² .	---

CAUTION:

Used only on above mentioned crops and follow the recommended safety period before harvest.

RE-ENTRY PERIOD:

24 hours after the last application.

COMPATIBILITY:

Compatible with most common used pesticides, incompatible with liquid fertilizers, it's recommended to mix **PROPORID** with water then add other pesticide before mixing.

• For more details about first aid & precautions please refer to first aid & precautions index.