

# FLUOXINIL-62.5% WDG

## FUNGICIDE

### WATER DISPERSIBLE GRANULES

**Class:**  
anilinopyrimidine  
phenylpyrrole

A systemic fungicide with protectant activity against a wide range of fungal diseases on agricultural crops mentioned in table of uses.

**COMPOSITION:** Each kg contains:

<b>CYPRONIL</b>	<b>37.5% W/W (A.I)</b>
<b>FLUDIOXONIL</b>	<b>25.0% W/W (A.I)</b>

**PROPERTIES:**

**FLUOXINIL-62.5%WDG:** a new fungicide used to control a broad spectrum of fungal disease and protect from resistant formulation towards other fungicide.

**FLUOXINIL-62.5% WDG:** Is a combination of two fungicides (CYPRONIL and FLUDIOXONIL) for the control of botrytis in Beans, Peas, Potato, Broccoli, Cabbage, Cauliflower, Kale, Mustard Greens, Cucumber, Watermelon, Melon, Pumpkin, Squash, Celery, Lettuce, Parsley, Spinach, Beet, Carrot, Radish, Turnip, Sweet Potato, Onion, Garlic, Tomatoes, Eggplant, Pepper, Okra, Avocado, Guava, Mango, Citrus, Grape, Strawberry

**FLUOXINIL-62.5%WDG:** Effective Fungicide with Long Residual Activity on Plant.

**USES AND RECOMMENDATIONS:**

CROPS	PESTS	APPLICATION RATE (GM/20 LITER)	SAFETY PERIOD (DAY)
Bean, Peas	White Mold, Gray Mold	15-20	7
Potato	Brown Rot, Early Blight, Septoria Leaf Spot, Powdery Mildew	15-20	14
Broccoli, Cabbage, Cauliflower, Kale, Mustard Greens.	Powdery Mildew	10 -17	7
	Alternaria Leaf Blight, Leaf Spot	15-20	
Cucumber, Watermelon, Melon, Pumpkin, Squash.	Alternaria Leaf Blight, Leaf Spot, Stem Blight, Powdery Mildew	15-20	7

<b>Celery, Lettuce, Parsley, Spinach</b>	Alternaria Leaf Blight, Leaf Spot, Mold, Powdery Mildew	15-20	7
<b>Beet, Carrot, Radish, Turnip, Sweet Potato</b>	Powdery Mildew , Alternaria Leaf Blight	15-20	7
<b>Onion, Garlic</b>	Leaf Blight, Purple Blotch, Neck Rot, Black Mold	15-20	7
	Soil Borne Disease: White Rot	10-20 (apply at the time of planting as an in furrow spray)	
<b>Tomato, Eggplant, Pepper, Okra</b>	Early Blight, Gray Mold, Powdery Mildew	15-20	7
<b>Avocado, Guava, Mango,</b>	Botrytis Fruit Mold, Leaf Spot, Anthracnose	15-20	21
<b>Lemon</b>	Stem End Rot, Anthracnose, Blue Mold, Gray Mold	15-20	21
<b>Grape</b>	Molds	15-20	14
<b>Strawberry</b>	Gray Mold, Powdery Mildew, Anthracnose	15-20	14

**POST HARVEST APPLICATIONS:**

<b>CROPS</b>	<b>DISEASE</b>	<b>APPLICATION RATE</b>	<b>REMARKS</b>
<b>Pomegranates</b>	Botrytis Fruit Rot, Gray Mold	29 Gm/ 1 Liter	Dip For Approximately 30 Seconds And Allow Fruit To Drain.

**CAUTION:**

Used Only On Above Mentioned Crops And Follow The Recommended Safety Period Before Harvest.

**COMPATIBILITY:**

Compatible with Many Commonly Used Fungicides and Insecticides but It's Preferable to Make a Compatibility Test before Mix.

**RE ENTRY PERIOD:**

24 Hours from the Last Spraying.