

ISO 9001 & 14001 REGISTERED



DNV Certification B.V., The Netherlands

**VAPCO**  
**VETERINARY & AGRICULTURAL**  
**PRODUCTS MANUFACTURING CO. LTD.**



**HANGER-50% WDG**  
**FUNGICIDE**  
**WATER DISPERSIBLE GRANULES**

**A SYSTEMIC FUNGICIDE USED TO CONTROL SCAB AND ROT AND POWDERY MILDEW ON CROPS MENTIONED IN THE TABLE OF USES**

**COMPOSITION:** Each kg contains:

**CYPRDONIL**

**50.0 % W/W (A.I)**

**PROPERTIES:**

**HANGER-50% WDG** : SYSTEMIC FUNGICIDE, UPTAKE INTO PLANTS AFTER FOLIAR APPLICATION AND TRANSPORT THROUGHOUT THE TISSUE AND ACROPETALLY IN THE XYLEM. INHIBITS PENETRATION AND MYCELIAL GROWTH BOTH INSIDE AND ON THE LEAF SURFACE.

**HANGER-50% WDG**: USED TO CONTROL SCAB ON APPLE, BLOSSOM BLIGHT AND BROWN ROT ON STONE FRUIT AND BOTRYTIS BUNCH ROT ON GRAPES.

**HANGER-50% WDG**: RAPIDLY DEGRADATION IN SOIL.



### USES AND APPLICATION RATE:

CROPS	DISEASE / TIMING	APPLICATION RATE (GM /20 LT WATER)	PHI (DAYS)	NOTES
APPLE	SCAB / (PREBLOOM STAGE )	11.25 (112.5 GM/200 LT WATER )	21	FIRST APPLICATION AT GREEN TIP
	SCAB / PREBLOOM, BLOOM POSTBLOOM STAGE.	6.0 (60 GM/200LT WATER) WHEN MIXED WITH (METIRAM OR MANCOZEB)		SECOND APPLICATION IS AFTER 7-10 DAYS FROM THE FIRST APPLICATION
	SCAB, POWDERY MILDEW/ PREBLOOM BLOOM, POSTBLOOM STAGE	6.0 (60 GM/200LT WATER) WHEN MIXED WITH (MYCLOBUTANIL)		
APRICOT, NECTARINES, PEACHS, PLUM AND PRUNES	BROWN ROT AND BLOSSOM BLIGHT / (BLOOM STAGE)	11.25 (112.5 GM/200 LT WATER )	2	SECOND APPLICATION IS AFTER 7-10 DAYS FROM THE FIRST APPLICATION
	FRUIT BROWN ROT / (LAST SEASON)	22.5 (225 GM/200LT WATER)  11.25 (112.5 GM/200 LT WATER) When mixed with (MYCLOBUTANIL)		APPLY 10-14 DAYS PRIOR TO HARVEST
GRAPES,	BOTRYTIS BUNCH ROT	22.5 (225 GM/200LT WATER)	7	BEGIN APPLICATION AT EARLY BLOOM ONLY IF WEATHER IS WET.  THE MOST IMPORTANT APPLICATION CAN BE AT EARLY RIPENING TO PRE-HARVEST

ISO 9001 & 14001 REGISTERED



MGMT. SYS.  
RvA C.024

MGMT. SYS.  
RvA C.425

DNV Certification B.V., The Netherlands

# VAPCO

## VETERINARY & AGRICULTURAL PRODUCTS MANUFACTURING CO. LTD.



### **NOTES:**

- USED ONLY ON APPLE STONE FRUITS AND GRAPES
- DO NOT APPLY ON CHERRIES
- VIGOROUS AGITATION IS NECESSARY FOR PROPER DISPERSAL OF THE PRODUCT
- IF TANK MIXED WITH WETTABLE POWDER, WETTABLE GRANULES, DRY FLOWABLE, LIQUID FLOWABLE LIQUID OR EMULSIFIABLE CONCENTRATE PRODUCT, PLEASE ADD HANGER-50% FIRST AND ALLOW IT TO BE COMPLETELY DISPERSED THEN ADD THE OTHER PRODUCT.

### **RE ENTRY PERIOD:**

- 72 HOURS AFTER THE LAST APPLICATION FOR APPLE AND STONE FRUIT ORCHARDS
- 48 HOURS AFTER THE LAST APPLICATION FOR GRAPES

### **CROP TOLERANCE:**

NON PHYTOTOXIC WHEN USED AS RECOMMENDED. DO NOT APPLY ON CHERRIES

### **COMPATIBILITY:**

SHOULD NOT BE MIXED WITH ANY PRODUCT WHICH PROHIBITS SUCH MIXING.