

METASTOR 39.1% SE

FUNGICIDE SUSPO-EMULSION

Class:
1- Methoxyacrylate
2. acylalanine

Preventative fungicide with systemic properties for control of many soilborne plant diseases.

COMPOSITION: Each liter contains:

AZOXYSTROBIN	28.2 % W/V (A.I)
METALAXYL-M	10.9% W/V (A.I)

PROPERTIES:

METASTOR 39.1%: Is a mixture of Azoxystrobin and Metalaxyl-m (Mefenoxam) compound used for the control of soil borne diseases on various vegetables and field crops.

METASTOR 39.1%: Used to control a wide range of fungal diseases such as rhizoctonia seedling blight (rhizoctonia solani), damping off (pythium), root rot on a wide range of agricultural crops which mentioned in table.

METASTOR 39.1%: Preventative fungicide with systemic properties to control a wide range of fungal disease.

USES AND APPLICATION RATES:

CROPS	PEST	APPLICATION RATE /20 LT OF WATER	SAFETY PERIOD
Garlic, Onion, Leeks	rhizoctonia seedling blight (rhizoctonia solani), damping off (pythium), root rot	13.2 ML (DRENCHING)	0 DAY
Muskmelon, Watermelon, Cucumber, Pumpkin, Squash	rhizoctonia seedling blight (rhizoctonia solani), damping off (pythium), root rot	13.2 ML (DRENCHING)	0 DAY
Celery, Lettuce, Parsley, Spinach, Swiss Chard	rhizoctonia seedling blight (rhizoctonia solani), damping off (pythium), root rot	13.2 ML (DRENCHING)	36 DAY
Bean, Chickpea, Lentil, Pea, Broad Bean	rhizoctonia seedling blight (rhizoctonia solani), damping off (pythium), root rot	13.2 ML (DRENCHING)	0 DAY
Beet, Radish, Sweet Potato, Turnip	rhizoctonia seedling blight (rhizoctonia solani), damping off (pythium), root rot	13.2 ML (DRENCHING)	0 DAY

- For more details about first aid & precautions please refer to first aid & precautions index.

CROPS	PEST	APPLICATION RATE /20 LT OF WATER	SAFETY PERIOD
Broccoli , Cabbage , Mustard Greens	rhizoctonia seedling blight (rhizoctonia solani), damping off (pythium), root rot	13.2 ML (DRENCHING))	36 DAY
Tomato, Pepper	rhizoctonia seedling blight (rhizoctonia solani), damping off (pythium), root rot, stem rot	20- 30 ML (DRENCHING)	0 DAY

NOTICE :

- Used only on above mentioned crops and follow the recommended safety period before harvest.
- The product is extremely phytotoxic to certain apple varieties, so do not spray uniform where spray drift may reach apples.
- Apply this product only one application per season.

RE-ENTRY PERIOD:

12 hours from the last spraying.

- For more details about first aid & precautions please refer to first aid & precautions index.