

PROXNIL-62.5% WDG

FUNGICIDE

WATER DISPERSIBLE GRANULES

Class:
 anilinopyrimidine
 phenylpyrrole

A systemic fungicide with protectant activity against a wide range of fungal diseases on agricultural crops mentioned in table of uses.

COMPOSITION: Each kg contains:

| | |
|--------------------|------------------------|
| CYPRDONIL | 37.5% W/W (A.I) |
| FLUDIOXONIL | 25.0% W/W (A.I) |

PROPERTIES:

PROXNIL-62.5% WDG: A New Fungicide Used To Control A Broad Spectrum Of Fungal Disease And Protect From Resistant Formulation Towards Other Fungicide.

PROXNIL-62.5% WDG: Is A Combination Of Two Fungicides (Cypridonil And Fludioxonil) For The Control Of Botrytis In Grapes, Bulb Onions, Botrytis, Anthracnose And Rhizopus In Strawberries And Botrytis And Anthracnose In Blueberries.

PROXNIL-62.5% WDG: Effective Fungicide with Long Residual Activity on Plant.

USES AND RECOMMENDATIONS:

| CROPS | PESTS | APPLICATION RATE (GM/20 LITER) | SAFETY PERIOD (DAY) |
|--|--|--------------------------------|---------------------|
| Beans | White Mold, Gray Mold | 15 - 20 | 7 |
| Potatoes | Brown Rot, Early Blight, Powdery Mildew, Septoria Leaf Spot, Tan Spot | 15 - 20 | 14 |
| Broccoli, Cabbage, Cauliflower, Kale, Mustard Greens | Powdery Mildew | 10 - 17 | 7 |
| | Alternaria Leaf Blight, Cercospora Leaf Spot | 15 - 20 | |
| Cucumber, Watermelon, Melon, Pumpkin, Squash. | Alternaria Leaf Blight, Septoria Leaf Spot, Gummy Stem Blight, Powdery Mildew. | 15 - 20 | 7 |

| | | | |
|---|---|---|----|
| Celery, Lettuce, Parsley, Spinach | Alternaria Leaf Spot, Septoria Leaf Spot, Gray Mold, Sclerotinia Rot, Basal Rot, Powdery Mildew | 15 - 20 | 7 |
| Beet, Carrot, Radish, Turnip, Sweet Potato | Powdery Mildew , Alternaria Leaf Blight | 15 - 20 | 7 |
| Onion, Garlic | Botrytis Leaf Blight, Purple Blotch, Neck Rot, Black Mold. | 15 - 20 | 7 |
| | Soil borne Disease: White Rot | 10 - 20 (applay at the time of planting as an in furrow spray) | |
| Tomatoes, Eggplant, Pepper, Okra | Early Blight, Grey Mold, Powdery Mildew, Target Spot | 15 – 20 | 7 |
| Avocado, Guava, Mango, | Botrytis Fruit Rot, Alternaria Leaf Spot, Anthracnose | 15 – 20 | 21 |
| Citrus | Alternaria Stem End Rot, Anthracnose, Blue Mold, Green Mold | 15 - 20 | 21 |
| Grapes | Grey Mold, Sour Rot (Caused By A Fungal Complex) | 15 - 20 | 14 |
| Strawberry | Gray Mold, Powdery Mildew, Anthracnose | 15 - 20 | 14 |

POST HARVEST APPLICATIONS:

| CROPS | DISEASE | APPLICATION RATE | REMARKS |
|---------------------|-------------------------------|-------------------------|--|
| Pomegranates | Botrytis Fruit Rot, Gray Mold | 29 Gm/ 1 Liter | Dip For Approximately 30 Seconds And Allow Fruit To Drain. |

CAUTION:

Used Only On Above Mentioned Crops And Follow The Recommended Safety Period Before Harvest.

COMPATIBILITY:

Compatible with Many Commonly Used Fungicides and Insecticides but It's Preferable to Make a Compatibility Test before Mix.

RE ENTRY PERIOD:

24 Hours from the Last Spraying.