



THEILACIDE

Injectable Solution
For Cattle

COMPOSITION: Each ml contains:

Buparvaquone 50 mg

INDICATIONS:

For the treatment of theileriosis (East Coast Fever, Corridor disease, Tropical theileriosis etc..) in cattle caused by theileria parva, T. annulata, T. mutans, and T. orientalis. It is active against both the Schizonts and piroplasm stages of theileria and may be used during the incubation period of the disease or when clinical signs are apparent.

DOSAGE AND ADMINISTRATION:

For intramuscular injection into the neck muscles.

Cattle: A single dose of 1ml per 20 kg bodyweight (equivalent to 2.5 mg activity buparvaquone/ kg bodyweight).

In severe cases, same dose might be repeated after 48-72 hrs after the initial injection (but into the other side of the neck).

Please Read Carefully The Below Remarks Before Usage:

- Localized, swelling may occasionally be seen at the injection site.
- Theileriosis has severe depressant effects on the immune system. Therefore it is recommended that any vaccination be delayed until the animal has recovered.
- Should never be given by intravenous or subcutaneous injection.
- Avoid wet or dirty areas of the skin.
- No more than 10 ml should be injected into a single site.

WITHDRAWAL PERIODS:

- For Meat consumption: 42 days
- For Milk consumption: 48 hours

STORAGE: Keep in dry, dark place at below 30 °C

For vet. use only