

# CYPOTHRIN-25%

## INSECTICIDE

### EMULSIFIABLE CONCENTRATE

**Class:**  
Pyrethriod

**A pyrethroidal insecticide effective against wide range of insect pests attacking agricultural crops.**

**COMPOSITION:** Each litre contains:

**CYPERMETHRIN**

**25% W/V (A.I)**

**PROPERTIES:**

**CYPOTHRIN-25%:** A pyrethroidal insecticide, moderately toxic to human and animals.

**CYPOTHRIN-25%:** Extremely active against cutworms, leafhoppers, flea beetles, cucumber beetles, aphid, whiteflies, armyworms, loopers, thrips on broccoli, cauliflower, cabbage, lettuce, garlic, leeks, and onion.

**CYPOTHRIN-25%:** Non phytotoxic when used as recommended.

**USES AND APPLICATION RATE:**

SAFETY PERIOD (DAYS)	APPLICATION RATE ML / 20 LITER WATER	PESTS	CROP
DAYS 7 2 - 8 weeks	10 - 5 10 - 7	Cutworms, tobacco budworm, bugs, caterpillar, leafhoppers, flea beetles, cucumber beetles, aphid, whiteflies.	<b>Broccoli, cauliflower, cabbage.</b>
DAYS 7	10 - 5 10 - 7	Armyworms, loopers, thrips Cucumber beetles, cutworms, diamondback moth, flea beetles, leafhoppers, tobacco budworm.	
DAYS 7	10 - 7	Armyworms, loopers, thrips	<b>Garlic, leeks, onion.</b>
DAYS 7	10 - 5	Thrips, aphid, armyworms, cutworms, leafminers, stink bugs	

**CAUTION:**

- Used only on above mentioned crops and follow the recommended safety period before harvest.
- Lower rate of **CYPOTHRIN-25%** should be used under light and moderate insect pressure.
- Higher rates of **CYPOTHRIN-25%** should be used to control heavy to extremely heavy insect populations.

- For more details about first aid & precautions please refer to first aid & precautions index.

**PHYTOTOXICITY:**

Not phytotoxic when used as recommended.

**RE-ENTRY PERIOD:**

48 hours from the last spraying.

**COMPATIBILITY:**

Compatible with most pesticides, but it's preferable to make compatible test before mix.

- For more details about first aid & precautions please refer to first aid & precautions index.