

LINCO – SPECTO

INJECTABLE SOLUTION

COMPOSITION:

Each ml Contains :

Lincomycin (Lincomycin HCl)	50 mg
Spectinomycin (Spectinomycin Sulphate)	100 mg

INDICATIONS:

Linco - Specto Injection is effective against infections caused by micro-organisms sensitive for lincomycin and spectinomycin.

It is indicated for the treatment of many appearing respiratory infections (with or without complications caused by Mycoplasmas) and gastrointestinal infections in calves, sheep, goats, poultry, dogs and cats; it's used for dysentery, foot rot in sheep and goats; CRD (with or without complications caused by E. coli) in poultry and turkeys.

DOSAGE AND ADMINISTRATION:

For intramuscular or subcutaneous injection.

- Calves, sheep, goats : 1 ml per 10 kg bodyweight daily, for 3 – 7 days.
- Dogs, cats : 1 ml per 5 kg bodyweight daily, for a maximum of 3days.
- Poultry, turkeys : 0.2 ml per kg bodyweight daily, for a maximum of 3 days.

CONTRA-INDICATIONS:

- Do not administer to animals hypersensitive to lincomycin or spectinomycin
- Do not administer to animals with blood disturbances or severe impaired liver function.
- Do not administer to rabbits, horses and ruminants with an active rumen and intestinal microbial flora.
- Do not use in layers producing eggs for human consumption or animals in lactation if milk is intended for human consumption.

SIDE EFFECTS:

Shortly after injection a light pain or itching can occur.

WITHDRAWAL PERIOD:

Meat : 14 days

Kidneys and liver : 21 days

STORAGE:

Store in a dry, dark place below 30 °C.

PACKING : 100 ml , 50 ml