

LINCO-SPECTO

Water Soluble Powder

COMPOSITION: Each 150 gm contains:

Lincomycin	33.3 gm
Spectinomycin	66.7 gm

INDICATIONS:

CHICKENS: For increase in rate of body weight gain, for prevention and treatment of chronic respiratory disease (airsacculitis) caused by Mycoplasma spp and CRD Complex (complicated airsacculitis) associated with mycoplasmal and coliform infections susceptible to lincomycin and/or spectinomycin.

TURKEYS: For use in turkey poultlets up to 5 days of age for prevention and treatment of airsacculitis caused by Mycoplasma meleagridis susceptible to lincomycin and/or spectinomycin.

DOSAGE:

• **CHICKENS**

*** FOR TREATMENT OF CHRONIC RESPIRATORY DISEASE (Airsacculitis) AND CRD COMPLEX (Complicated Airsacculitis):**

Administer in drinking water at a rate of 0.5 g antibiotic activity per liter. Provide as the sole source of water for three days. Begin treatment at the first signs of disease.

*** FOR INCREASE IN RATE OF BODY WEIGHT GAIN:**

Administer 50 mg of antibiotic activity per kg of body weight per day with drinking water during the first five to seven days of life.

*** FOR PREVENTION OF CHRONIC RESPIRATORY DISEASE (Airsacculitis) AND CRD COMPLEX (Complicated Airsacculitis):**

Administer 150 mg of antibiotic activity per kg of body weight per day with drinking water for the first three to five days of life and 50 mg of antibiotic activity per kg of body weight per day for one to two days at about four weeks of age or at the time of vaccination.

• **TURKEYS**

FOR TREATMENT AND PREVENTION OF AIRSACCULITIS

Administer 150 mg of antibiotic activity per kg of body weight per day with drinking water for the first three to five days of life.

VAPCO

ANTIBACTERIALS/
ANTIBIOTICS

WITHDRAWAL PERIOD:

For meat consumption: 2 days before slaughtering.

WARNING:

- If no improvement is seen within three to five days, re-determine the diagnosis.
- Do not allow species other than chickens and turkeys access to the medicated drinking water.

STORAGE:

Keep in a dry dark place below 30 °C.

Packing: 150 gm.