Melodone

Injectable solution

Composition: Each ml contains:

Meloxicam 20 mg

Indictions:

For cattle : for use in acute respiratory infection with appropriate antibiotic therapy to reduce clinical signs , In diarrhea in combination with oral re- hydration therapy to reduce clinical signs in calves of over one week of age and young, non- lactating cattle , for adjunctive therapy in the treatment of acute mastitis , in combination with antibiotic therapy.

For horses: for use in the alleviation of inflammation and relief of pain in both acute and chronic musculoskeletal disorders, for the relief of pain associated with equine colic.

Dosage:

CATTLE: single subcutaneous or intravenous injection. 0.5 mg meloxicam / kg body weight (2.5 ml /100 kg body weight) in combination with antibiotic therapy or with oral re- hydration therapy as appropriate.

HORSE: single intravenous injection

0.6 mg meloxicam / kg body weight (3 ml / 100 kg body weight). melodone 15 mg / ml oral suspension may be used for continuation of treatment at a dosage of 0.6 mg meloxicam / kg body weight, 24 hours after administration of the injection.

Contraindications:

- Do not use in the following cases:
- In horses less than 6 weeks of age.
- In cattle less than one week of age for treatment of diarrhea.
- In pregnant or lactating mares.

-



- In animals suffering from impaired hepatic, cardiac or renal functions and hemorrhagic disorders, or where there is evidence of ulcerogenic gastrointestinal lesions.
- In case of hypersensitivity to the active substance or to any of the excipients.

Remarks:

- In cattle, a transient swelling at the injection site can occur but resolves without intervention.
- In horses, anaphylactic reactions can occur and should be treated symptomatically.
- Do not administer concurrently with gluco- corticosteroids, other non steroidal anti- inflammatory drugs or with anti- coagulant agents

Withdrawal periods:

Cattle: for meat consumption: 15 day For milk consumption: 5 days

Horses: for meat consumption: 5 days

Storage: keep in dry place at below 30 °C.

PACKING: 50 ML, 100 ML.