

VAPCO

INJECTABLE
PRODUCT

SPIROJECT

Injectable Solution

COMPOSITION:

Each ml contains:

Spiramycin 600 000 IU

PROPERTIES:

- Spiramycin is an antibiotic of the macrolides group . It is bacteriostatic and bactericidal at high concentrations. Its mode of action is similar to that of macrolides: it inhibits peptide synthesis by dissociating the RNA complex during translocation.
- Spiramycin is effective against Gram-positive bacteria and mycoplasmas, as well as certain Gram-negative germs, particularly Legionella and Actinobacillus.
- It spectrum of activity includes Staphylococcus spp., Streptococcus spp., Enterococcus, Streptococcus pneumoniae, Neisseria intracellularis, Listeria monocytogenes, Corynebacterium diphtheriae, Mycoplasma spp., Clostridium spp., Bacteroides spp., Haemophilus spp., Treponema pallidum, Treponema microdentium, Leptospira icterohemorrhagiae, Toxoplasma, Miyagawanella, Brucella, Pasteurella, Rickettsia spp. and Amoeba spp.

INDICATIONS:

Cattle : Prevention and the treatment of pulmonary infections, gastroenteritis, mastitis, metritis, omphalitis and omphalophlebitis, arthritis and foot -rot.

POSOLOGY AND METHOD OF ADMINISTRATION

Intramuscular route

Calves: 5 ml of SPIROJECT/ 40 kg b.w. as single injection or 2 administrations at 24 hours interval (75 000 IU of Spiramycin/kg b.w.)

Cattle: 5 ml of SPIROJECT/ 100 kg b.w. as single injection or 2 administrations at 24 hours interval (30 000 IU of Spiramycin/kg b.w.)

VAPCO

INJECTABLE
PRODUCT

Do not inject more than 15 mL per injection site.

CONTRA-INDICATIONS

Do not treat animals hypersensitive to macrolides.

WITHDRAWAL PERIODS

- Meat: 21 days
- Milk: 7 days or 14 milkings

STORAGE

Keep in dry place below 30 ° C.

PACKING: 50 ML, 100 ML.