



# **FLUAZIVAP-15% EC**

## **HERBICIDE**

### **EMULSIFIABLE CONCENTRATE**

#### **INTRODUCTION:**

A SELECTIVE POST-EMERGENCE HERBICIDE FOR CONTROL OF ANNUAL AND PERENNIAL GRASSES MENTIONED IN TABLE OF USES.

#### **COMPOSITION:** EACH LITER CONTAINS:

**FLUAZIFOP-P-BUTYL** **15.0 % W/V (A.I)**

#### **PROPERTIES:**

**FLUAZIVAP – 15% EC** : SYSTEMIC HERBICIDE TO CONTROL MANY ANNUAL AND PERENNIAL GRASS WEEDS IN MOST BROADLEAF CROPS MENTIONED IN THE TABLE OF USES.

**FLUAZIVAP – 15% EC:** ABSORBED MAINLY THROUGH LEAVES AND IS RECOMMENDED FOR USE AFTER CROP AND WEED EMERGENCE.

**FLUAZIVAP – 15% EC:** SELECTIVE, SYSTEMIC HERBICIDE, ABSORBED BY THE LEAVES AND TRANSLOCATED BOTH ACROPETALLY AND BASIPETALLY IN THE PHLOEM AND XYLEM, ACCUMULATING IN MERISTEMS.



# VAPCO

## VETERINARY & AGRICULTURAL PRODUCTS MANUFACTURING CO. LTD.



### USES AND RECOMMENDATIONS:

CROP	WEEDS CONTROLLED	RATE	CRITICAL COMMENTS
<p><b>ASPARAGUS, BEANS, CAPSICUMS, CARROTS, SQUASH, PUMPKIN, WATERMELON, CUCUMBER, MUSKMELON LUCERNE, ONIONS, PEAS, POTATOES, TOMATOES, BROCCOLI, CABBAGE, BRUSSELS SPROUT, CAULIFLOWER, COLLARD GREEN, KALE, TURNIP, RADISH.</b></p>	<p>ANNUAL GRASSES AND PERENNIAL GRASS REGROWTH</p>		<p>ASPARAGUS: APPLY AFTER THE FINAL PICK. DO NOT APPLY DURING THE CUTTING SEASON. OPTIMAL CONTROL WILL BE OBTAINED BY TREATING ANNUAL GRASSES AT THE SEEDLING STAGE AND PERENNIAL REGROWTHS AT THE 4 TO 5 LEAF STAGE.</p>
<p><b>BERRYFRUIT, GRAPVINES, KIWIFRUIT, APPLE, PEAR, PEACHES, PLUM, CHERRIES, NECTARINE, APRICOT, OLIVE</b></p>	<p>ANNUAL SEEDLING GRASSES AND PERENNIAL GRASS REGROWTH</p>	<p>KINDLY CHECK THE BELOW SPRAYING TABLE.</p>	<p>NON-BEARING FRUIT CROPS: MAY BE APPLIED SAFELY OVER THE CROP (SEE CAUTION BELOW). BEARING FRUIT CROPS: SHOULD BE APPLIED AS A DIRECTED SPRAY TO AVOID CONTACT WITH THE FRUIT. CAUTION: APPLICATION SHOULD NOT BE MADE OVER FRUIT CROPS UNDER CONDITIONS OF HIGH TEMPERATURE (25°C) AND HIGH LIGHT INTENSITY AS MINOR LEAF SCORCH MAY OCCUR ON SENSITIVE SPECIES SUCH AS PEACHES.</p>
<p><b>ORNAMENTALS (SEEDLING AND ESTABLISHED),</b></p>	<p>ANNUAL SEEDLING GRASSES AND PERENNIAL GRASS REGROWTH</p>		<p>MAY BE USED SAFELY OVER THE TOP OF A WIDE RANGE OF ORNAMENTALS AND SHELTER TREES INCLUDING FLAX, BULBS AND CABBAGE TREES. DO NOT APPLY TO PAMPAS GRASS, BAMBOO AND OTHER ORNAMENTAL GRASSES AND TUSsockS. CAUTION: APPLICATION SHOULD NOT BE MADE OVER ORNAMENTALS UNDER CONDITIONS OF HIGH TEMPERATURE (25°C) AND HIGH LIGHT INTENSITY AS MINOR LEAF SCORCH MAY OCCUR ON SENSITIVE SPECIES.</p>
<p><b>WHITE CLOVER (SEED CROPS)</b></p>	<p>ANNUAL GRASSES AND PERENNIAL GRASS REGROWTH</p>		<p>APPLY IN MID OCTOBER AFTER CLOSING UP. REMOVE STOCK AND ALLOW THE GRASS TO FRESHEN UP WITH 5 TO 7 CM OF NEW LEAF, BEFORE MAKING TREATMENT. MCPB CAN BE TANK MIXED WITH FLUAZIVAP FOR BROADLEAF WEED CONTROL.</p>



- **SPRAYING TABLE:**

GRASS SPECIES	GROWTH STAGE		
	SEEDLING	VEGETATIVE	MATURE
BARNYARD GRASS	1.25 TO 1.7 LITRES	2.5 LITRES	NOT RECOMMENDED
BRISTLE GRASS	1.25 TO 1.7 LITRES	2.5 LITRES	NOT RECOMMENDED
CANARY GRASS	1.7 LITRES	1.7 LITRES	NOT RECOMMENDED
COUCH (TWITCH)*	1.7 LITRES	1.7 LITRES	NOT RECOMMENDED
CROWFOOT GRASS	1.7 LITRES	1.7 LITRES	1.7 LITRES
INDIAN DOAB	2.5 LITRES	2.5 LITRES	2.5 LITRES
MERCER GRASS	1.7 LITRES	2.5 LITRES	2.5 LITRES
PASPALUM	1.7 LITRES	2.5 LITRES	2.5 LITRES
PRAIRIE GRASS	0.85 LITRES	0.85 LITRES	1.7 LITRES
RYEGRASSES (EXCLUDING PERENNIAL SPECIES)	1.7 LITRES	2.5 LITRES	NOT RECOMMENDED
SOFT BROME (GOOSE GRASS)	1.7 LITRES	NOT RECOMMENDED	NOT RECOMMENDED
SUMMER GRASS	1.25 TO 1.7 LITRES	NOT RECOMMENDED	NOT RECOMMENDED
VOLUNTEER CEREALS	0.85 LITRES	0.85 LITRES	NOT RECOMMENDED
WILD OATS	0.85 LITRES	0.85 LITRES	NOT RECOMMENDED

**RE-ENTRY PERIOD:**

DON'T ENTER FIELD BEFORE 48 HOURS FROM LAST APPLICATION.

**CROP TOLERANCE:**

NOT PHYTOTOXIC WHEN USED AS RECOMMENDED.

**COMPATIBILITY:**

COMPATIBLE WITH MOST PRODUCTS, BUT IT'S PREFERABLE TO MAKE COMPATIBLE TEST BEFORE MIX.