



## **FLY BAIT-0.06% GR**

### **PUBLIC HEALTH INSECTICIDE**

### **GRANULES**

**INTRODUCTION:**

**FLY BAIT AIDS IN AID** IN THE REDUCTION OF HOUSEFLY POPULATIONS AROUND INDUSTRIAL BUILDINGS, COMMERCIAL FACILITIES, AND AGRICULTURAL STRUCTURES AND PREMISES

**COMPOSITION:** EACH KG CONTAINS:

**METAFLUMIZONE                      0.06 % W/W (A.I)**

**PROPERTIES:**

**FLY BAIT-0.06% GR:** IS INSECTICIDAL BAIT WHICH AIDS IN THE REDUCTION OF COMMON HOUSEFLY (*MUSCA DOMESTICA*) POPULATIONS

**FLY BAIT-0.06% GR:** NON-SYSTEMIC PUBLIC HEALTH INSECTICIDE PRIMARILY ACTIVE BY INGESTION.

**FLY BAIT-0.06% GR:** CONTAINS THE FLY ATTRACTING PHEROMONE, Z-9 TRICOSENE

**FLY BAIT-0.06% GR:** CAN BE BROADCAST OVER LARGER AREAS AS A SCATTER BAIT OR IT CAN BE PLACED IN BAIT STATIONS/TRAYS



### USES AND RECOMMENDATIONS:

APPLICATION SITES	APPLICA TION METHOD	APPLICATI ON TIMING	APPLIC ATION RATE	APPLICATION DIRECTIONS
<p>AROUND THE OUTSIDE OF FOOD AND BEVERAGE PROCESSING PLANTS, MEAT AND POULTRY PROCESSING PLANTS, FOOD HANDLING ESTABLISHMENTS, RESTAURANTS, CAFES, FAST-FOOD ESTABLISHMENTS, SUPERMARKETS, FARM MARKETS, BAKERIES, WAREHOUSES, FACILITIES RECREATIONAL AREAS INCLUDING PARKS, PICNIC GROUNDS, CAMP GROUNDS, AND OUTDOOR LATRINES, LIVESTOCK HANDLING AND FEEDING FACILITIES INCLUDING BROILER AND LAYER HOUSES, LIVESTOCK BARNs, HORSE STABLES, FEED STORAGE BUILDINGS, FEED SILOS, AND OTHER ANIMAL HANDLING AND FEEDING FACILITIES.</p>	SCATTER BAIT	APPLY FLY BAIT WHEN FLIES ARE OBSERVED , TO KEEP POPULATIO NS AT MANAGEAB LE LEVELS.	2.5 G/ M <sup>2</sup>	SCATTER FLY BAIT GRANULES EVENLY ONTO DRY, LEVEL SURFACES WHERE FLIES ARE OBSERVED TO BE RESTING AND FEEDING. DO NOT PLACE BAIT IN PILES OR IN WET/MOIST AREAS. FOR OUTDOOR USE, DO NOT REAPPLY THE PRODUCT MORE THAN 3 WEEKS APART.
INDOORS IN LIVESTOCK HANDLING AND FEEDING FACILITIES, BUT ONLY IN AREAS WHERE LIVESTOCK CANNOT COME IN CONTACT WITH OR INGEST BAIT GRANULES				
<p>AROUND AREAS INFESTED WITH HOUSEFLIES INCLUDING FOOD AND BEVERAGE PROCESSING PLANTS, MEAT AND POULTRY PROCESSING PLANTS AND FOOD HANDLING INSTITUTIONS, RESTAURANTS, CAFES, FAST FOOD ESTABLISHMENTS, SUPERMARKETS, FARM MARKETS, BAKERIES, WAREHOUSES, LIVESTOCK HANDLING AND FEEDING FACILITIES INCLUDING MEAT AND LAYERED HOMES, PIG PRODUCTION FACILITIES, HORSE STABLES, DAIRY SHEDS, FEED STORAGE BUILDINGS, FEED SILOS</p>	BAIT STATION S/TRAYS	APPLY FLY BAIT AT THE BEGINNING OF THE SEASON TO PREVENT LARGE INFESTATI ONS AND TO KEEP POPULATIO NS AT MANAGEAB LE LEVELS.	28-56 G	USE FLY BAIT IN ANY COMMERCIAL FLY BAIT STATION PLACED AT LEAST 1.2 M ABOVE THE GROUND. DO NOT PILE OR HEAP BAIT IN STATION OR TRAY. TO INSURE ADEQUATE COVERAGE, HANG OR PLACE 8 TO 10 STATIONS PER 180 M2 OF FLOOR SPACE IN AREAS WHERE FLIES ARE OBSERVED TO BE RESTING OR FEEDING. REAPPLY AS BAIT IS CONSUMED.

### USE RESTRICTIONS

ISO 9001 & 14001 REGISTERED



# VAPCO

## VETERINARY & AGRICULTURAL PRODUCTS MANUFACTURING CO. LTD.



- DO NOT APPLY **FLY BAIT** INSIDE HOMES.
- APPLY **FLY BAIT** ONLY IN LOCATIONS WHERE CHILDREN AND PETS CANNOT ACCESS OR COME INTO CONTACT WITH BAIT GRANULES.
- DO NOT APPLY THIS PRODUCT IN A WAY THAT WILL CONTACT WORKERS OR OTHER PERSONS, EITHER DIRECTLY OR THROUGH DRIFT.
- DO NOT APPLY IN ANY AREA WHERE LIVESTOCK CAN INGEST OR CONTACT THE BAIT.
- EXCEPT FOR BAIT STATION USE, DO NOT USE IN EDIBLE PRODUCT AREAS OF FOOD HANDLING ESTABLISHMENTS, RESTAURANTS,
- OR OTHER AREAS WHERE FOOD IS COMMERCIALY PREPARED OR PROCESSED OR IN SERVING AREAS WHILE FOOD IS EXPOSED.
- DO NOT APPLY THIS PRODUCT WHEN RAINFALL IS FORECAST WITHIN 48 HOURS.
- DO NOT MAKE BROADCAST APPLICATION WITHIN 4.5 M OF PERENNIAL FRESH WATER BODIES, INCLUDING LAKES, RESERVOIRS, RIVERS, PERMANENT STREAMS, MARSHES, NATURAL PONDS, AND COMMERCIAL FISHPONDS.
- DO NOT MAKE BROADCAST APPLICATION WITHIN 18 M OF ESTUARINE WATER BODIES (BRACKISH WATER), INCLUDING TIDAL WATER SUCH AS BAYS, MOUTHS OF RIVERS, SALT MARSHES, AND LAGOONS.

### **RE-ENTRY PERIOD:**

USED AS PUBLIC HEALTH INSECTICIDE

### **COMPATIBILITY:**

INCOMPATIBLE WITH MOST PESTICIDES.