

TRIPLE SULPHA Advance

Water Dispersible Powder
For Poultry, Ruminants

COMPOSITION: Each gm contains

Sulphadimidine Na	250 mg
Sulphadiazine Na	250 mg
sulfamethoxazole	250 mg

PROPERTIES:

- 3-Combined valuable sulpha components, with a high antibacterial activity against Gm +ve and Gm -ve bacteria: Coli, Staph, Strept, Salmonella, Pasteurella, Proteus, Chlamydia etc.
- Also effective against Eimeria spp. (Coccidia).
- Sulphonamides diffuse very widely into tissues and thus exert effectiveness against the bacterial infections.
- Sulphonamides are well tolerated and have low toxicity; hence they are safe to be used.

INDICATIONS:

- **Poultry:** Coryza, Caecal coccidiosis.
- **Ruminants:** Pneumonia, Enteritis, Scour, Foot rot, Haemorrhagic septicaemia (Pasteurella)

DOSAGE:

Mix well before use.

- **Treatment:** Duration for 3-5 days.

Species	TRIPLE SULPHA advance
Poultry	250 gm per 200 liters of drinking water
Ruminants	10-15 gm per 50 kg B. wt. daily and orally with water

- **Prevention:** Half dosage for 3 days.

WITHDRAWAL PERIOD:

- For meat consumption: 7 days.
- For milk consumption: 4 days.

REMARKS:

- Not recommended for young stocks and layers.
- Ensure adequate water intake during treatment to reduce the risk of crystalluria.

STORAGE: Keep in a dry, dark place below 30 °C.

For vet. Use only