



CYPERMETHRIN-10 EC

Pyrethroidal Insecticide
Emulsifiable Concentrate

COMPOSITION : Each ml contains :

Cypermethrin **100 mg**

PROPERTIES :

- Non-systemic insecticide with contact and stomach action.
- It is formulated for use as dip or spray.
- Well tolerated by farm animals.
- Biodegradable does not accumulate in the tissues or in milk.
- Has rapid knockdown effects on all ticks including those resistant to organo-phosphorus and organo-chlorine insecticides.
- Very active at low concentration and has low mammalian toxicity.

USES:

- For the treatment of scab and mange mites of livestock beside poultry mites as well.
- For the control of different ectoparasites (ticks, fleas, lice, keds and screw worms) attacking cattle, sheep, goats and camels.
- For the control of nuisance flies and other insects in farms, stables and animals houses.

METHODS OF APPLICATION:

By spraying or dipping after dilution with water as follows:

Species	Dilution Rate	Treatment may be repeated after 1-2 weeks
Cattle, Camels	1:1000	
Flies on cattle	1:300	
Sheep, goats	1:800	
Poultry	1:750 spray (60 liters per 1000 birds)	
Animals Houses	1:50	



VAPCO

**VETERINARY & AGRICULTURAL
PRODUCTS MANUFACTURING CO. LTD.**



PRECAUTIONS:

- Toxic to fishes and bees.
- Avoid application on sick, weak and young animals.
- Avoid contact with eyes, mouth, skin and inhalation.
- Before spraying, wear protective clothes, boots, gloves and goggles.
- Do not eat, smoke or drink during spraying.
- Avoid polluting ponds, waterways and water sources.
- Do not allow off-springs to suckle until mothers are completely dried from the insecticide liquid.

FIRST – AID:

- Eye contact: wash with water for 15 minutes.
- Skin contact: wash with soap and water.
- Inhalation:
 - Move patient to fresh air
 - Apply artificial respiration, give oxygen when necessary.
- Ingestion:
 - Do not induce vomiting.
 - Apply gastric lavage.
- Get medical help, when necessary.

ANTIDOTE:

- Not specific, treat according to symptoms.
- Give diazepam in case of convulsions.

WITHDRAWAL PERIOD: For meat and milk consumption: One week.

STORAGE:

- Keep away from children, food and drink.
- Keep in a dry, dark place below 30 °C.

For vet use only