



DELTAVAP-5% EC

Pyrethroidal Insecticide
Emulsifiable Concentrate

COMPOSITION: Each ml contains:

Deltamethrin **50 mg**

PROPERTIES:

- Non-systemic insecticide with contact and stomach action. Powerful and fast acting.

USES:

Treatment of mange and for the control of ticks, mites, keds, lice, fleas and other insects attacking sheep, goats, cattle, camels, and their premises.

METHODS OF APPLICATIONS:

By spraying or dipping after dilution with water as follows:

	DIP	SPRAY
Ticks	1:2000 Replenish 1:700	10 ml / 20 liters (1:2000)
Mange	1:1000 Replenish 1:400	20 ml / 20 liters (1:1000)
Lice	1:4000 Replenish 1:1200	5 ml / 20 liters (1:4000)
Flies	1:2000 Replenish 1:700	10 ml / 20 liters (1:2000)
Animals houses		75 ml / 20 liters (1:250)

PRECAUTIONS:

- Toxic to fishes and bees.
- Avoid application on sick, weak and young animals
- Avoid contact with eyes, mouth, skin and inhalation.
- Before spraying, wear protective clothes, glove, boots, and goggles.
- Do not eat, drink or smoke during application.
- Do not allow off-springs to suckle until mothers are completely dried from the insecticide liquid.
- Avoid polluting ponds, water ways and water sources.



VAPCO

**VETERINARY & AGRICULTURAL
PRODUCTS MANUFACTURING CO. LTD.**



FIRST – AID:

- Eye contact: wash with water for 15 minutes.
- Skin contact: wash with soap and water.
- Inhalation:
 - Move patient to fresh air
 - Apply artificial respiration, give oxygen when necessary.
- Ingestion:
 - Do not induce vomiting.
 - Apply gastric lavage.
- Get medical help, when necessary.

ANTIDOTE:

- Not specific, treat according to symptoms.
- Give diazepam in case of convulsions.

WITHDRAWAL PERIOD:

For meat and milk consumption: 2 days.

STORAGE:

- Keep away from children, food and drinks.
- Keep in a dry, dark place below 30 °C.

For vet use only