



DIAZINON-25 EC

Organophosphorus Insecticide And Acaricide
Emulsifiable Concentrate

COMPOSITION : Each ml contains :

Diazinon **250 mg**

PROPERTIES :

- Non-systemic insecticide and acaricide with contact, stomach and respiratory action.
- A powerful and broad spectrum insecticide and acaricide.
- Well tolerated by farm animals and pets except cats and poultry.

USES:

- Treatment of scab, mange and for the control of mites, ticks, lice, keds, fleas and screw worms on sheep, goats, cattle and camels.
- Control of stable flies and blow-fly larvae.

METHODS OF APPLICATIONS: By spraying or dipping after dilution with water as follows:

Species	spray	dip
Sheep, goats,	1:400	1:1000 Replenish 1:350
Cattle, Camels	1:400	1:400 Replenish 1:160
Animals houses	1:10	-----

PRECAUTIONS:

- Diazinon must be used directly after being diluted with water, and not kept for the next day to be used especially for cattle.
- In presence of water, the stored diazinon is oxidized to diazoxon which is harmful to animals especially cattle.
- Opened bottles should be used within 3 months.
- Do not use after expiration date.
- Toxic to fishes, bees, cats, poultry and birds.
- Avoid application on sick, exhausted, weak and young animals.
- Before spraying, wear protective clothes, boots, gloves and goggles.
- Do not eat, drink or smoke during spraying.
- Avoid polluting ponds, waterways and water sources.
- Do not allow off-springs to suckle until mothers are completely dried from the insecticide liquid.



VAPCO

**VETERINARY & AGRICULTURAL
PRODUCTS MANUFACTURING CO. LTD.**



FIRST – AID:

- Eye contact: wash with water for 15 minutes.
- Skin contact: wash with soap and water.
- Inhalation:
 - Move patient to fresh air
 - Apply artificial respiration, give oxygen when necessary.
- **Ingestion:**
 - Do not induce vomiting.
 - Apply gastric lavage.
- Get medical help, when necessary.

ANTIDOTE:

- Atropine, PAM, Toxogonin.
- Continue symptomatic treatment.

WITHDRAWAL PERIOD:

- For meat consumption: 14 days.
- For milk consumption: 3 days.

STORAGE:

- Keep away from children, food and drinks.
- Keep in a dry, dark place below 30 °C.

For vet use only