



DUSTRIN-5%

Ready – To – Use
 Dusting Powder Insecticide

COMPOSITION: Each gm contains:

Malathion **50 mg**

PROPERTIES:

- Non-systemic insecticide and acaricide with contact, stomach and respiratory action.
- Organo-phosphorus ectoparasiticide with the highest efficacy and lower toxicity.

USES:

- Eradication of ectoparasites (Lice, Fleas, Ticks, Keds, Sarcoptic mange mites and horn flies) infesting poultry, cattle, sheep, goats, camels, equines, rabbits, dogs, cats and game-birds (Quails, partridges etc...).
- Control of wound-infesting maggots which attack the animal.
- Combat of flies, mosquitoes, cockroaches, bugs etc..., in and around farm buildings, different animals housing and institutional uses.

METHOD OF APPLICATION: (Directly, Ready – To – Use Insecticide).

- Dust thoroughly (with avoidance of the eyes) over the body, the ears, between the folds of the skin, under the wings and around the vent (for birds); then rub with hand between wool / hairs / fur / feathers. (15 gm / bird, 50 gm/ animal).
- Milking animals: apply only after milking, then wash the udder before the next milking.
- Avoid contamination of watering and feeding troughs, eggs, equipment, and utensils.
- Repeat application every two weeks if needed.
- Stables, poultry houses (Cages, nests, roosts of birds), pens, kennels, cat's sleeping baskets, houses and institutional uses: should thoroughly be dusted, with more attention to corners and cracks where parasites hide.
(75-125 gm / sq. meter of litter or floor space.

PRECAUTIONS:

- Toxic to bees and fishes.
- Avoid contact with eyes, mouth, skin and inhalation.
- Wear protective clothes, gloves, boots and face shield before dusting.
- Do not eat, drink or smoke during dusting.
- Do not treat sick, exhausted, weak or very young animals.
- Avoid polluting water-sources, ponds and water canals.
- Always, dispose empty containers safely.



VAPCO

**VETERINARY & AGRICULTURAL
PRODUCTS MANUFACTURING CO. LTD.**



FIRST – AID :

■ **Ingestion :**

- Induce vomiting by throat finger, touch only.
- Apply gastric lavage
- Eyes contact: Flush with water for 15 minutes.
- Inhalation:
 - Apply artificial respiration, and give oxygen, if necessary.
 - move patient to fresh air.
- Skin contact: Wash with soap and water.
- Get medical help, when necessary

ANTIDOTE:

- Atropine, PAM.
- continue symptomatic treatment.

WITHDRAWAL PERIOD:

For meat consumption: 7 days.

STORAGE:

- Keep away from children, food and drink.
- Keep in a dry, dark place below 30 °C.

For vet use only