



# VAPCO

**VETERINARY & AGRICULTURAL  
PRODUCTS MANUFACTURING CO. LTD.**



# VAP MALATHION-50% EC

**Insecticide and Acaricide  
Emulsifiable Concentrate**

**COMPOSITION:** Each ml contains:

**Malathion**

**500 mg**

**PROPERTIES:**

- Non-systemic insecticide and acaricide with contact, stomach and respiratory action.
- Strong organo-phosphorus ectoparasiticide which is effective at low concentration.
- Low mammalian toxicity.

**USES:**

- Treatment of different types of mange.
- Control of many species of ticks, mites, lice, keds, fleas, flies, mosquitoes, attacking livestock.
- Control of wound-infesting maggots which attack the animal.
- For the combat of insects in farms, stables and other animals houses.

**METHODS OF APPLICATION:**

| Species        | Dilution Rate | application   |
|----------------|---------------|---------------|
| Cattle         | 1:100         | Spray         |
| Sheep, Goats   | 1:100         | Spray or dips |
| pigs           | 1:100         | Spray         |
| Equine         | 1:100         | Spray         |
| Poultry        | 1:200         | Spray         |
| Animals Houses | 1: 25         | Spray         |

**PRECAUTIONS:**

- Toxic to bees and fishes.
- Avoid contact with eyes, mouth, skin and inhalation.
- Wear protective clothes, gloves, boots and face shield before spraying.
- Do not eat, drink or smoke during spraying.
- Do not treat sick, exhausted, weak or very young animals.
- Avoid polluting water-sources, ponds and water canals.
- Remove lactating animals & all animals under 1 month old before application.



# VAPCO

**VETERINARY & AGRICULTURAL  
PRODUCTS MANUFACTURING CO. LTD.**



## REMARKS:

- Repeat at 2-3 weeks interval, if necessary.
- For horses use: 1.9 liters of diluted liquid per animal.
- For horses: use 40 – 80 liters of the diluted liquid per 1000 sq. meter.
- For poultry houses: clean then spray floors, walls, ceiling, roosts and nests.
- For poultry: use 38 liters of 0.5% spray solution per 1000 birds.

## FIRST – AID:

- Eye contact: wash with water for 15 minutes.
- Skin contact: wash with soap and water.
- Inhalation:
  - Move patient to fresh air
  - Apply artificial respiration, give oxygen when necessary.
- Ingestion:
  - Do not induce vomiting.
  - Apply gastric lavage.

Get medical help, when necessary

## ANTIDOTE:

- Atropine, PAM, Toxogonin.
- continue symptomatic treatment.

**WITHDRAWAL PERIOD:** For milk consumption: one day.

## STORAGE:

- Keep away from children, food and drink.
- Keep in a dry, dark place below 30 °C.

**For vet use only**