



VAPCOCIDIN- 20 EC

Pyrethroidal Insecticide and Acaricide
Emulsifiable Concentrate

COMPOSITION: Each ml contains:

Fenvalerate **200 mg**

PROPERTIES:

- Non-systemic insecticide and acaricide with contact and stomach action.
- Controls a wide range of external parasites, including those resistant to other insecticides.
- High insecticidal, ovocidal and larvicidal activity at low dosage.
- It is characterized by low toxicity to mammals.
- Has a powerful antecedent and repellent effects.

USES:

- For the treatment of different forms of mange and for the control of ticks, lice and keds on sheep, goats, cattle and camels.
- It is used to combat flies on cattle, stable flies, and crawling insects in shelters, animals and poultry houses.
- Is sprayed on fly breeding sites to destroy larvae and keep down fly population.

METHODS OF APPLICATIONS:

By spraying or dipping after dilution with water as follows:

USE	DILUTION	Interval
Dip	1:2500 Replenish. 1:1300	Weekly or as required
Spray	100 ml per 20 liters of water (1:200)	
For stables, and animals houses	1:100	

PRECAUTIONS:

- Avoid contact with eyes, mouth, skin, and inhalation.
- Don't allow off-spring to suckle until mothers are completely dried from the insecticide liquid.
- Don't mix with other insecticides.
- Toxic to bees and fishes.
- Before spraying, wear protective clothes, gloves, boots, and goggles.
- Do not eat, drink or smoke during spraying.
- Avoid polluting ponds, waterways and water sources.
- Don't treat sick, exhausted, weak or very young animals.



VAPCO

**VETERINARY & AGRICULTURAL
PRODUCTS MANUFACTURING CO. LTD.**



FIRST – AID:

- Eye contact: wash with water for 15 minutes.
- Skin contact: wash with soap and water.
- Inhalation:
 - Move patient to fresh air
 - Apply artificial respiration, give oxygen when necessary.
- Ingestion:
 - Do not induce vomiting.
 - Apply gastric lavage.
- Get medical help, when necessary.

ANTIDOTE:

- Not specific, treat according to symptoms.
- Give diazepam in case of convulsion.

WITHDRAWAL PERIOD:

For meat consumption: 2 days.

STORAGE:

- Keep in a dry, dark place below 30 °C.
- Keep away from children, food and drinks.

For vet use only