



VAPCOPOUR

Suspension Concentrate Insecticide

COMPOSITION: Each ml contains:

Imidacloprid 350 mg

PROPERTIES:

- Systemic insecticide with broad – spectrum effect against external parasite (fleas, lice, ticks.....etc)
- Has a contact and stomach action

INDICATIONS:

- Control of fleas, lice & ticks attacking sheep, goats, cattle, camels, dogs and cats; and their premises.

METHOD OF APPLICATION:

- Topical use: pour 1 ml per animal along the back area with the contact of skin.
- Large animals (cattle and camels) may need more than (1 ml)
- For dogs and cats apply on the back between the shoulders where it can not be licked of, to avoid any oral toxicity. According to the doses in the table below.

Animal	Weight	Dose
Cats	Up to 4 kg	0.15 ml
	Over 4 kg	0.25 ml
Dogs	Up to 4 kg	0.15 ml
	4 – 10 kg	0.3 ml
	10 – 25 kg	0.7 ml

- ✓ Do not use in kittens less than 8 weeks or puppies less than 7 weeks of age.
- ✓ One application lasts for one month.

Premises:

Soak 3 pieces of sponges (3 × 5 cm size each) with the product and distribute carefully in the stable (house) in different sites, which are away from and difficult to be accessed by children and animals.

FIRST – AID:

- Eye – contact: wash with water for 15 min.
- Inhalation: move patient to fresh air, and artificial respiration is applied when necessary.
- Ingestion: move the patient to hospital.
- Anti – Dote: not specific treat according to symptoms.

ISO 9001 & 14001 REGISTERED



DNV Certification B.V., The Netherlands

VAPCO

**VETERINARY & AGRICULTURAL
PRODUCTS MANUFACTURING CO. LTD.**



PRECAUTIONS:

- Toxic to bees
- Toxic to fishes
- Before use, wear protective clothes, gloves and goggles.
- Avoid contact with eyes, mouth, skin and inhalation.
- Avoid polluting water sources, ponds and water canals.
- Dispose empty containers safely.

STORAGE:

- Keep in dry dark place below 30°C.
- Keep away from children, food and drinks.

For vet use only