



VAPCOZIN – 12.5 EC

Insecticide and Acaricide
Emulsifiable Concentrate

COMPOSITION: Each ml contains:

Amitraz **125 mg**

PROPERTIES:

- Non-systemic insecticide and acaricide of the amidine group with contact and respiratory action.
- Characterized by its strong effect against ticks which are resistant to arsenical, organo-phosphorus and organo-chlorine insecticides.
- Rapidly degraded in the soil and does not cause environmental pollution hazards.
- Low mammalian toxicity.
- Expellent action causes ticks to withdraw mouth parts rapidly and fall off the host animal.

USES:

- Treatment of different strains of scab and mange.
- Control of many species of ticks, mites, lice, keds, fleas and flies on sheep, goats, cattle, camels and dogs.
- For combating insects in farms, stables and other animals houses.
- Destruction of flies-larvae in breeding sites and keeping down flies population.

METHOD OF APPLICATIONS:

By spraying or dipping after dilution with water as follows:

Species	Initial dilution rate	Interval	Replenishment
Cattle, Camels: Ticks, Lice Mange	1:700 1:500	7-10 days 12 days	Replenishment rate is half of the initial dilution rate
Sheep, Goats Ticks Scab Keds, Lice	1:700 1:250 1:250	At need At need 10-12 days	
Dogs: Ticks, Mange	1:250	10 days	
Animals Houses	2-4 %	At need	



VAPCO

VETERINARY & AGRICULTURAL PRODUCTS MANUFACTURING CO. LTD.



PRECAUTIONS:

- Toxic to equines and fishes.
- Avoid contact with eyes, mouth, skin and inhalation.
- Before spraying, wear protective clothes, gloves, boots and goggles.
- Do not eat, drink or smoke during spraying.
- Avoid polluting ponds, water ways and water sources.
- Amitraz should not be used in animals with known hypersensitivity or allergy to the drugs
- Amitraz should be avoided in animals less than four months of age
- Caution must be used in diabetic animals
- Caution must be used when applying this to very small toy breeding dogs.
- Amitraz may interact with corticosteroids and azathioprine
- Do not allow off-spring to suckle until mothers are completely dried from the insecticide liquid.

FIRST – AID:

- Eye contact: wash with water for 15 minutes.
- Skin contact: wash with soap and water.
- Inhalation:
 - Move patient to fresh air
 - Apply artificial respiration, give oxygen when necessary.
- Ingestion:
 - Do not induce vomiting.
 - wash the mouth with water.
 - let patient, if conscious , to drink 1-2 glasses of water to dilute the product in the stomach.
 - Apply gastric lavage.

Get medical help, when necessary

ANTIDOTE: Not specific, treat according to symptoms.

WITHDRAWAL PERIOD:

- For milk consumption: 2 days.
- For meat consumption: Sheep ,Goats: 7 days.
Cattle: 1 day.

STORAGE:

- Keep away from children, food and drinks.
- Keep in a dry, dark place below 30 °C.

For vet use only