



AMOXAN

Water Soluble Powder

COMPOSITION: Each gm contains:
Amoxycillin Trihydrate
200 mg

PROPERTIES:

- A beta-lactam antibiotic of the aminopenicillins.
- Safe, less toxic and rapidly absorbed from the gastrointestinal tract.
- Acts by impairing development of bacterial cell walls.
- Has a bactericidal action against a wide range of Gm+ve and Gm-ve bacteria, attacking chickens, pigeons, ducks, pre-ruminating animals, dogs and cats:
 Staph, Strept, Coryne, Listeria, Clostridia, Erysipelothrix; E. Coli, Salmonella, Pasteurella, Moraxella, Klebsiella, Proteus, Haemophilus, Shigella, Actinobacillus, Leptospira, Borrelia, Spirochaeta.

INDICATIONS:

- CRD, Colibacillosis, Salmonellosis, Fowl cholera, Bovine haemorrhagic septicaemia, Clostridial infections, Omphalitis and yolk sac infection, Coryza, Sinusitis, Air succulitis, Diarrhea, Enteritis, Spirochaetosis, Listeriosis, Erysipelas.
- Other respiratory, gastrointestinal and urogenital infections.
- Stress conditions due to vaccination, debeaking, transporting, rehousing, changes of feed and weather.

DOSAGE: (Firstly, dissolve well in 2 liters of water, then add to the water bulk to be medicated).

- **Treatment:** (Duration: 5 days)
 - **Poultry:** 100 gm / 200 liters of drinking water.
 - **Lambs, Calves, kids** (Pre-ruminating):
 2 gm / 50 kg B. wt. twice daily and orally with drinking water.
 (8 mg activity / kg body wt.)
 - **Dogs, Cats:**
 0.5 gm / 10 kg B. wt. twice daily and orally with drinking water.
 (10 mg activity / kg body wt.)
 - **Pigeons, Ducks:**
 0.5 gm / 5 kg B. wt. with drinking water.
 (20 mg activity / kg body wt.)
- **Prevention:** Half of the dosage for the same period.

WITHDRAWAL PERIOD: (For Meat Consumption)

- For Poultry, Pigeon, Ducks: 3 Days.
- For Lambs, Kids, Calves: 8 Days



VAPCO
VETERINARY & AGRICULTURAL
PRODUCTS MANUFACTURING CO. LTD.



■ **REMARKS:**

- Don't use in poultry producing eggs for human consumption.
- Not used for rabbits.

STORAGE: Keep in dry, dark place below 30 °C.

PACKING: (100, 250, 500, 1000) GM.

For Vet. Use only