



# AMOXYCOL

Water Soluble Powder

For Poultry, Cattle, Sheep, Goats, Calves, Foals, Horses

**COMPOSTION:** Each gm contains:

Amoxycillin Trihydrate	200 mg
Colistin Sulphate	800,000 I.U.

## PROPERTIES:

- A combination of two wide – spectrum antibiotics which leads to a strong and synergistic action against poultry and livestock diseases.
- **AMOXYCOL** is used in a large-scale variety of infections of respiratory, alimentary, and urogenital tracts caused by Gm + ve and Gm – ve pathogenic bacteria such as: Coli, Salmonella Clostridia, Mycoplasma, Pasteurella, Haemophilus, Proteus, Actinomyces, Staphylococcus, Streptococcus, Coryne, Listeria, Yersinia, Moraxella, Bordetella, Actinobacillus, Klebsiella etc....

## INDICATIONS:

- Salmonellosis, Colibacillosis, Clostridial infections, CRD, Omphalitis and Yolk sac infection, Sinusitis, Coryza, Air Succulitis, Pasteurellosis, Mycoplasmosis, Leptospirosis.
- Cases of enteritis, septicemia, secondary bacterial and generalized infections.
- Stress conditions due to vaccination, debeaking, transporting, rehousing, changes of feed and weather.

**DOSAGE:** (Firstly, dissolve well in 2 liters of water, then add to the water bulk to be medicated).

- **Treatment:** (for 3-5 days)
  - **Poultry:** 100 gm per 200 liters of drinking water.
  - **Cattle, Horses:** 8-10 gm per animal twice daily and orally with water.
  - **Calves, Foals:** 3-6 gm per animal twice daily and orally with water.
  - **Sheep, Goats:** 2-5 gm per animal twice daily and orally with water.
- **Prevention:** Half dosage for the same period.

## WITHDRAWAL PERIOD:

- For meat consumption: 1 week.

## REMARKS:

- Not used for layers.
- Do not use in lactating animals
- Not used for rabbits.

**STORAGE:** Keep in dry, dark place below 30 °C.

**PACKING:** (100, 250, 500, 1000) GM.

**For Vet. Use only**