



AMPIVAP-50%

Water Soluble Powder
For Poultry, calves, lambs and kids

COMPOSITION: Each gm contains:

Ampicillin Trihydrate **500 mg**

DESCRIPTION:

Is a semi-synthetic penicillin with a broad spectrum activity against a large number of Gram-negative and some Gram-positive bacteria. Acts bactericidal by interfering with the cell wall synthesis of bacteria.

INDICATIONS:

- Is indicated for the treatment of enteritis in calves, lambs and kids caused by *E. coli*, pneumonia caused by *Pasteurella multocida*, and infections caused by *Salmonella dublin*.
- In poultry, is effective in the treatment of enteritis and airway infections caused by *E. coli*, *Salmonella* and *Clostridium spp.*
- is indicated for the treatment of dermatitis, caused by *Streptococcus spp.*; the treatment of enteritis, caused by *E. coli*, *Campylobacter sputorum* and *Clostridium perfringens* type C; the treatment of urogenital infections caused by *Corynebacterium spp.*

DOSAGE:

- For oral administration via drinking water or milk replacer. Rinse the drinking water systems thoroughly before and after administration of the product.

- **Calves, lambs, kids:** 20 – 40 mg per kg bodyweight, twice daily, during 5 days.
- **Poultry:** Use amount of the product as stated in table in 1,000 litres of drinking water.

Age in days – Grams of product

0 – 7	250
8 – 14	305
15 – 21	363
22 – 28	421
29 – 35	486
36 – 42	557

- Medicated drinking water should be used within 4 hours.

CONTRA-INDICATION:

- Do not administer to adult ruminants.
- Do not administer to animals with an impaired renal function.
- Do not use for treatment of infections caused by penicillinase producing micro-organisms.
- Do not administer to animals hypersensitive to penicillins.
- Do not use in layers producing eggs for human consumption.

O 9001 & 14001 REGISTERED



VAPCO

VETERINARY & AGRICULTURAL PRODUCTS MANUFACTURING CO. LTD.



SIDE-EFFECTS:

Sometimes gastro-intestinal disturbances, hypersensitivity reactions to penicillins or allergic skin reactions may occur.

INCOMPATIBILITY WITH OTHER DRUGS:

Do not combine with bacteriostatic antimicrobial drugs.

WITHDRAWAL PERIOD:

Meat:

Calves: 2 days, Poultry: 1 day.

STORAGE: Keep in dry, dark place below 30 °C.

PACKING: (100, 250, 500, 1000) GM.

For Vet Use Only