



# VAZURIL 5%

(Suspension)

For Calves and Lambs

**COMPOSITION: Each 1 ml contains:**

**Toltrazuril**

**50 mg**

## Indication:

- Lambs: The prevention of clinical signs of coccidiosis and reduction of coccidian shedding in Lambs on farms with a confirmed history of coccidiosis caused by *Eimeria crandallis* and *Eimeria ovinoidalis*.
- Calves: For the prevention of clinical signs of coccidiosis and reduction of coccidia shedding in housed calves replacing cows producing milk for human consumption (dairy cows) on farms with a confirmed history of coccidiosis caused by *Eimeria bovis* or *Eimeria zuernii*.

**Dosage:** Shake well before use.

- Lambs: Single oral dose of 0.4 ml oral suspension per kg body weight (20 mg toltrazuril/kg body weight).
- Calves: Single oral dose 3.0 ml oral suspension per 10 kg body weight (corresponding to 15 mg toltrazuril/kg body weight).
- 

## Withdrawal period:

- Calves: 63 days, not permitted for use in lactating animals producing milk for human consumption.
- Lambs: Meat and offal: 42 days, not permitted for use in lactating sheep producing milk for human consumption.

## Contraindications:

- Do not use in calves weighing more than 80 kg bodyweight.
- Do not use in fattening units such as veal or beef calves.

ISO 9001 & 14001 REGISTERED



**VAPCO**  
**VETERINARY & AGRICULTURAL**  
**PRODUCTS MANUFACTURING CO. LTD.**



**Precautions:**

- Frequent and repeated use of antiprotozoal from the same class may lead to the development of resistance.
- Avoid contact with skin, and dispose the unused drug.

**Storage:**

Keep in dry, dark place below 30°C.

**For vet use only**