

FLY BAIT

PUBLIC HEALTH INSECTICIDE
GRANULES

VAPCO

Class:
Semicarbazone

A public health insecticide that aids in the reduction of housefly populations around industrial buildings, commercial facilities, and agricultural structures and premises

COMPOSITION: Each kg contains:

METAFLUMIZONE

0.06% W/W (A.I)

PROPERTIES:

FLY BAIT: Is insecticidal bait which aids in the reduction of common housefly (*Musca domestica*) populations.

FLY BAIT: Non-systemic public health insecticide primarily active by ingestion.

FLY BAIT: Contains the fly attracting pheromone, Z-9 tricosene.

FLY BAIT: Can be broadcast over larger areas as a scatter bait or it can be placed in bait stations or trays.

USES AND APPLICATION RATE:

APPLICATION SITES	APPLICATION METHOD	APPLICATION TIMING	APPLICATION RATE	APPLICATION DIRECTIONS
Outdoors: Around the outside of food and beverage processing plants, meat and poultry processing plants, restaurants, cafes, fast-food establishments, supermarkets, farm markets, bakeries, warehouses, facilities recreational areas including parks, picnic grounds, camp grounds, and outdoor latrines, livestock handling and feeding facilities including broiler and layer houses, livestock barns, horse stables, feed storage buildings, feed silos, and other animal handling and feeding facilities.	Scatter bait	Apply when flies are observed	2.5 g/ m ²	Scatter FLY BAIT granules evenly onto dry surfaces where flies are observed to be resting and feeding. Do not place bait in piles or in moist areas. For outdoor use, do not reapply the product more than 3 consecutive weeks apart.
Indoors: livestock handling and feeding facilities, but only in areas where livestock cannot come in contact with or ingest bait granules	Scatter bait		2.5 g/ m ²	

• For more details about first aid & precautions please refer to first aid & precautions index.

APPLICATION SITES	APPLICATION METHOD	APPLICATION TIMING	APPLICATION RATE	APPLICATION DIRECTIONS
Indoors: Around areas infested with Houseflies including food and beverage processing plants, Meat and poultry processing plants, Restaurants, cafes, fast food establishments, supermarkets, farm markets, bakeries, warehouses, livestock handling and feeding facilities including meat and layered homes, pig production facilities, horse stables, dairy sheds, feed storage buildings, feed silos, and other animal feeding facilities.	Bait stations /trays	Apply at the beginning of the season to prevent large infestations	28-56 g /m ²	Use FLY BAIT in any commercial fly bait stations or trays specialized for house flies. Station placed at least 1.2 m above the ground. Do not pile or heap bait in station or tray. To ensure enough coverage, hang or place 8 to 10 stations per 180 m ² of floor space. Reapply as bait is consumed.

CAUTION:

- Do not apply **FLY BAIT** inside homes.
- Apply **FLY BAIT** only in locations where children and pets cannot access or come into contact with the product granules.
- Do not apply this product in a way that will contact workers or other persons.
- Do not apply in any area where livestock can ingest or contact the bait.
- Do not apply the product by scattering, in areas of food handling establishments, restaurants, or other areas where food is commercially prepared or processed
- Do not apply this product when rainfall is forecast within 48 hours.
- Do not make broadcast application within 4.5 m of water bodies, including lakes, rivers, ponds, and commercial fishponds.
- Do not make broadcast application within 18 m of ground water.

RESIDUAL ACTIVITY: Has long residual activity effect that remains for more than 3 months

• For more details about first aid & precautions please refer to first aid & precautions index.