

# BENOMYL-50%WP

FUNGICIDE  
WETTABLE POWDER

VAPCO

Class:  
Benzimidazole

A systemic fungicide effective against a wide range of fungal diseases which affect agricultural crops.

**COMPOSITION:** Each kg contains:

**BENOMYL**

**50% W/W (A.I)**

## PROPERTIES:

**BENOMYL-50% WP:** A systemic fungicide for the control of fungal diseases which affect agricultural and horticultural crops and have a secondary effect on spidermites eggs.

**BENOMYL-50% WP:** Highly efficient at low concentration of use.

**BENOMYL-50% WP:** Harmless to humans, animals and bees.

**BENOMYL-50% WP:** Effective to keep the products after harvest and during storage without any side effect.

## USES AND APPLICATION RATE:

CROP	DISEASE	APPLICATION RATE/ 20 L WATER	TIME OF APPLICATION	SAFETY PERIOD (DAYS)
Apple, Pear, Quince, Almond, Apricot	Powdery mildew, Scab, Brown rot, Cherry leaf spot, Monilia	12 gm	Start spraying when first symptoms of disease appear, apply again at 10-14 days interval	14
				3
Grapes	Powdery mildew	12 gm	Before flowering apply again at 10-14 days interval	7
Potatoes	Powdery mildew	8 gm	Begin spraying when first symptoms of disease appear, repeat spray every 10-14 days if necessary	14
Tomatoes	Soft rot, Leaf rot, White rot, Leaf spots	5-6 gm	----	3
Squash, Cucumber, Watermelon, Muskmelon	Anthracnose, Downy mildew, Leaf spots	12 gm	Begin spraying when first symptoms of disease appear, repeat spray every 10-14 days	3
Strawberries	Powdery mildew, Leaf spot, Wilt, Root rot	20 gm	Start spraying when first symptoms of disease appear, apply again every two weeks. Cover booms for 25 minutes with <b>BENOMYL-50% WP</b> solution.	3

• For more details about first aid & precautions please refer to first aid & precautions index.

CROP	DISEASE	APPLICATION RATE/ 20 L WATER	TIME OF APPLICATION	SAFETY PERIOD (DAYS)
Bananas	Banana Rot	16 gm	After picking, dip the bananas in <b>BENOMYL-50% WP</b> for 10 minutes, then leave it to dry.	---
Beans	Powdery mildew	12 gm	Begins spraying when first symptoms of disease appear, repeat spray every 10-14 days.	14
Roses and ornamentals	Powdery mildew, Scab, Leaf spot, Grey mold	12 gm	Re-spray every 10-14 days interval if necessary.	21

**CAUTION:**

Used only on the above mentioned crops and follow the recommended safety period before harvest.

**REMARKS:**

Mix **BENOMYL-50% WP** with sticking material (Such as **SUPER SURFIX**) to ensure the uniform distribution on plants like Roses.

**RE-ENTRY PERIOD:**

24 hour from last application.

**COMPATIBILITY:**

Compatible with most insecticide and fungicide, but compatibility test is recommended before spraying.

**NOTICE:**

Humidity may cause chemical changes of **BENOMYL-50% WP**, therefore, the container should be closed tightly and kept away from humidity.

• For more details about first aid & precautions please refer to first aid & precautions index.