# **COPROXIDE-77% WP**

## FUNGICIDE WETTABLE POWDER



A protective and curiative fungicide on a wide range of agricultural crops.

**COMPOSITION:** Each kg contains:

COPPER HYDROXIDE

77% W/W (A.I)

(EQV. 50% W/W COPPER (A.I))

PROPERTIES:

COPROXIDE-77% WP: Is a contact fungicide and bactericide with a preventive action against

a wide range of diseases.

COPROXIDE-77% WP: Control bacterial infections on agricultural crops such as Angular leaf

spot.

#### **USES AND APPLICATION RATE:**

CROP	DISEASE	APPLICATION RATE GM / 20 LT. WATER	SAFETY PERIOD (DAYS)
Squash	Anthracnose, Angular Spots, Downy Mildew	17 - 35	7
Potato	Early and Late Blight	10 - 45	15
Tomatoes	Bacterial Spot, Early and Late Blight, Anthracnose	20 - 45	10
Eggplant	Anthracnose, Blight	20	10
Almond, Apricot, Plum, Peach, Cherry	Rot, Shot Hole, Leaf Curl	35 - 50	15
Apple, Pear	Anthracnose, Black Spot, Fire Blight.	50 - 70	15
Orange, Lemon, Mandarin, Grapefruit	Black Rot.	45 - 90	21
Olives	Peacock Spot	35 - 40	21

<sup>·</sup> For more details about first aid & precautions please refer to first aid & precautions index.

### CAUTION:

- Used only on the above mentioned crops and follow the recommended safety period before harvest.
- Not phytotoxic for treated plants except some species of apples (Red apple), but may cause damage to plum and peach leaves especially at low temperature.

### NOTE:

Used inside plastic houses on tomatoes, pepper and eggplant.

### **RE-ENTRY PERIOD:**

48 hours from last application.

### **COMPATIBILITY:**

Compatible with most other pesticide except acidic compounds or Calcium sulphate and compatibility test is recommended before 24 hours from application

<sup>·</sup> For more details about first aid & precautions please refer to first aid & precautions index.