

DIFLUNAM- 40% SC

Fungicide
Suspension Concentrate

VAPCO

Class:
Cinnamamide⁽¹⁾
Dinitroaniline⁽²⁾

Foliar fungicide with protectant and translaminar activity, for control fungal disease listed in the table of uses

COMPOSITION: Each liter contains:

Dimethomorph ⁽¹⁾	20% W/V (A.I)
Fluazinam ⁽²⁾	20% W/V (A.I)

PROPERTIES:

DIFLUNAM-40% SC : A foliar fungicide, with protectant and translaminar activity, it is applied before the disease appears and is followed by a programme of repeat sprays

DIFLUNAM-40% SC : Used to control many fungal diseases to the crops mentioned in the table of use

USES AND APPLICATION RATE:

CROP	PESTS	DOSE	PHI
Potato	Late blight, Mold	75 ml/100 L water	7 days
Vegetables	Mold	75 ml/100 L water	7 days
Grape	Mold	75 ml/100 L water	14 days
Melon	Downy Mildew	50 ml/100 L water	7 days

CAUTION:

- Maximum number of application/season: 5.
- Do not apply by any type of irrigation system.
- Time of application:
* First Application Timing: Apply as soon as conditions favor disease incidence, or when disease warning is given, but before any disease appears.

Second and Repeat Applications Timings

Low Disease Risk: Spray at 14 day intervals.

High Disease Risk: Spray at 10 day intervals.

COMPATIBILITY:

Compatible with other pesticide and foliar fertilizers, compatibility test is recommended before spraying.

• For more details about first aid & precautions please refer to first aid & precautions index.