

DODIVAP- 40.7% SC

Fungicide
Suspension Concentrate

VAPCO

Class:
Guanidine

Systemic fungicide used to control various fungal diseases

COMPOSITION: Each liter contains:

Dodine

40.7% W/V (A.I)

PROPERTIES:

DODIVAP-40.7% SC: Is a systemic fungicide with protectant and some curative activity

DODIVAP-40.7% SC: Is a fungicide used to control various fungal diseases in the crops mentioned in the table of uses

USES AND APPLICATION RATE :

CROP	DISEASE	DOSE (/20 L WATER)	APPLICATION DIRECTIONS:	GENERAL REMARKS	NO. OF APPLICATION /SEASON	TIME BETWEEN APPLICATION
Almonds, chestnuts	Brown rot (<i>Monilinia fructicola</i>)	71 ml	Used at pink bud, full bloom, and petal fall	- Do not apply later than 5 weeks after petal fall.	3 days	10 days
	Scab (<i>Venturia carpophila anamorph</i>)	71 ml	Used when Petal fall, repeat application 2-5 weeks later.			
	Shot hole (<i>Wilsonomyces carpophilus</i>)	71 ml	used when leaf open until the petals fall, repeat application 2-5 weeks later.			
	Anthracnose (<i>Gnomonia leptostyla</i>)	71 ml	Used when first leaves unfold. Repeat application at 10- to 14-day intervals			
Apple	Scab (<i>Venturia inaequalis</i>)	35 ml	Used at the beginning at green tip, apply preventively at weekly intervals from green tip to petal fall.	- Do not apply after petal fall.	2 days	7 days

- For more details about first aid & precautions please refer to first aid & precautions index.

CROP	DISEASE	DOSE (/20 L WATER)	APPLICATION DIRECTIONS:	GENERAL REMARKS	NO. OF APPLICATION /SEASON	TIME BETWEEN APPLICATION
Apricots, Nectarines and Peaches	Leaf Curl (<i>Taphrina deformans</i>)	71 ml	Used at dormant or delayed dormant application. Additional applications may be made just before buds swell in the spring.	- Do not apply after petal fall.	3 days	7 days
	Brown rot (<i>Monilinia fructicola</i>)	71 ml	Used at pink bud. Repeat application at full bloom and again at petal fall			
	Scab (<i>Venturia carpophila anamorph</i>)	71 ml	Used at petal fall.			
	Shot hole (<i>Wilsonomyces carpophilus</i>)	71 ml	Used at dormant state, and repeat application at petal fall.			
Bananas	Sigatoka (<i>leaf spot</i>)	20-36 ml	-Air application	---	5 days	7-15 days
Cashews, Filberts, Walnuts	Brown rot (<i>Monilinia fructicola</i>)	71 ml	Used at pink bud, full bloom, and petal fall	- Do not apply within 3 days of harvest	3 days	10-14 days
	Scab (<i>Venturia inaequalis</i>)	71 ml	Used at petal fall, Repeat application 2-5 weeks later.			
	Shot hole (<i>Wilsonomyces carpophilus</i>)	71 ml	Used at leaf emergency (bloom through petal fall). Repeat application 3 and 5 weeks later			
	Anthraxnose (<i>Gnomonia leptostyla</i>)	71 ml	Used when first leaves unfold. Repeat application at 10- to 14-day intervals			
Cherries	Leaf Spot (<i>Blumeriella jaapii anamorph</i>)	35-71 ml	Used at petal fall or when first leaves unfold. Repeat after at 7- to 10-day intervals.	- Do not apply when temperatures and humidity are excessively high, because may cause spotting to foliage or fruit.	2 days	7-10 days
	Brown rot (<i>Monilinia fructicola</i>)	71 ml	Used at the time flower buds begin to open (white bud-popcorn bloom stages). Repeat application at full bloom and again at petal fall.	- Do not apply within 7 days of harvest.		

CROP	DISEASE	DOSE (/20 L WATER)	APPLICATION DIRECTIONS:	GENERAL REMARKS	NO. OF APPLICATION /SEASON	TIME BETWEEN APPLICATION
Peanuts	Early Leaf Spot (<i>Cercospora arachidicola</i>) and Late Leaf Spot (<i>Cercosporidium personatum</i>)	21-35 ml	Apply at begin with applications, first approximately 35 days after planting, and repeat at 10- to 14-day intervals.	- Do not apply within 14 days of harvest. - Do not feed hay or vines (peanuts) to livestock.	5 days	10-14 days
Pears	Scab (<i>Venturia pinna</i>) Leaf spot (<i>Fabraea spp.</i>)	71 ml	Apply at beginning at green tip	- Do not apply within 7 days of harvest.	3 days	7 days
Plums	Leaf Curl (<i>Taphrina deformans</i>)	71 ml	Apply dormant or delayed dormant application. Additional applications may be made just before buds swell in the spring.	- Do not apply within 7 days of harvest.	3 days	7 days
	Brown rot (<i>Monilinia fructicola</i>)	71 ml	Apply at pink bud. Repeat application at full bloom and again at petal fall			
	Scab (<i>Venturia carpophila anamorph</i>)	71 ml	Used at petal fall. Repeat application at 2 and 5 weeks after petal fall.			
	Shot hole (<i>Wilsonomyces carophilus</i>)	71 ml	Apply at dormant stage, and repeat application at petal fall.			

CAUTION:

- Used only on above mentioned crops.
- Do not apply through any type of irrigation system.
- Do not apply in greenhouses.
- Do not graze livestock in treated fields, orchards.

COMPATIBILITY:

Compatible with insecticides, fungicides. Herbicide and Acaricide, incompatible with some copper, oil, sulfur, foliar nutrients and Chlorpyrifos formulation, compatibility test is recommended before spraying.

RE ENTRY PERIOD:

48 hours from the last application