

# FLARE-50% WDG

FUNGICIDE  
WATER DISPERSIBLE GRANULES

VAPCO

Class:  
Hydroxylanilide

Non-systemic, foliar fungicide with protective activity. Inhibits germ tube elongation and mycelium growth to control of botrytis and monilinia diseases in many crops listed in the table of uses and use to suppress the powdery mildew disease in grapes.

COMPOSITION: Each Kg contains:

FENHEXAMID

50% W/W (A.I)

#### PROPERTIES:

**FLARE-50% WDG:** non-systemic protective fungicide used to control gray mold and blossom blight and twig blight (monilinia spp.) on many fruit trees and vegetables listed in the table of uses.

#### USES AND APPLICATION RATE:

CROP	DISEASE	APPLICATION RATE/20 LITER WATER	APPLICATION TIME	REMARKS	SAFETY PERIOD (DAYS)
Almonds	Blossom blight, twig blight (monilinia spp.)	34 gm	<ul style="list-style-type: none"><li>• Pink bud</li><li>• Full bloom</li><li>• Petal fall</li><li>• Up to 28 days after petal fall</li></ul>	<ul style="list-style-type: none"><li>- Apply in sufficient water to obtain complete coverage of flowers, fruit and twigs</li><li>- Do not make more than two consecutive applications with 7-14 days between applications</li></ul>	3
Cucumber	Gray mold (botrytis cinerea)	34 gm	<ul style="list-style-type: none"><li>• Begin applications when conditions favor disease development</li><li>• Repeat at 7-10 day if necessary</li></ul>	<ul style="list-style-type: none"><li>- Do not apply in field.</li><li>- Do not make more than 2 consecutive applications</li></ul>	7
Eggplant, pepper	Gray mold (botrytis cinerea)	34 gm	<ul style="list-style-type: none"><li>• Begin applications when conditions favor disease development</li><li>• Repeat at 7-10 day if necessary</li></ul>	<ul style="list-style-type: none"><li>- Do not apply in field.</li><li>- Do not make more than 2 consecutive applications</li></ul>	7

• For more details about first aid & precautions please refer to first aid & precautions index.

CROP	DISEASE	APPLICATION RATE/20 LITER WATER	APPLICATION TIME	REMARKS	SAFETY PERIOD (DAYS)
Tomato	Gray mold (botrytis cinerea)	34 gm	<ul style="list-style-type: none"> <li>Begin applications when conditions favor disease development</li> <li>Repeat at 7-10 day if necessary.</li> </ul>	<ul style="list-style-type: none"> <li>- Do not apply in field.</li> <li>- Do not make more than 2 consecutive applications.</li> </ul>	3
Lettuce, parsley	Gray mold (botrytis cinerea)	22- 34 gm	<ul style="list-style-type: none"> <li>Begin applications when conditions favor disease development</li> <li>Repeat at 7-10 day if necessary.</li> </ul>	<ul style="list-style-type: none"> <li>- Do not apply in field.</li> <li>- Do not make more than 2 consecutive applications.</li> </ul>	3
Grapes	Botrytis bunch rot (botrytis cinerea) Powdery mildew Suppression (uncinula necator)	22 gm	<ul style="list-style-type: none"> <li>• Early bloom.</li> <li>• Bunch pre-closure.</li> <li>• Veraison.</li> <li>• Up to day of harvest.</li> </ul>	<ul style="list-style-type: none"> <li>- Fenhexamid has been shown to enhance the efficacy of other fungicides when used in a powdery mildew spray program.</li> <li>- Do not make more than 2 consecutive applications with 7 days between applications</li> </ul>	3
Kiwi	Botrytis fruit rot (botrytis cinerea)	2 gm	Post-harvest	<ul style="list-style-type: none"> <li>- Do not make more than one application to fruit</li> <li>- Use only as a post-harvest application</li> <li>- (fruit dip treatment) Fruit should remain in dip solution for 20 to 30 seconds</li> </ul>	---
Strawberries	Gray mold (bolry/is cinerea)	34 gm	<ul style="list-style-type: none"> <li>• Begin application at early bloom.</li> <li>• Repeat at 7-10 day if necessary.</li> </ul>	<ul style="list-style-type: none"> <li>- Do not make more than 2 consecutive applications</li> </ul>	7

• For more details about first aid & precautions please refer to first aid & precautions index.

CROP	DISEASE	APPLICATION RATE/20 LITER WATER	APPLICATION TIME	REMARKS	SAFETY PERIOD (DAYS)
Apricot, cherries, nectarines, peaches.	Brown rot, blossom blight, twig blight (monilinia spp.)	34 gm	<ul style="list-style-type: none"> <li>• Early bloom.</li> <li>• Late bloom.</li> <li>• 2 weeks after petal fall.</li> <li>• Up to day of harvest.</li> </ul>	- Do not make more than 2 consecutive applications	3
	Gray mold on cherries (botrytis cinerea)	22 – 34 gm	<ul style="list-style-type: none"> <li>• Early bloom.</li> <li>• Repeat at 7-10 day if necessary.</li> <li>• Up to day of harvest.</li> </ul>		

**CAUTION:**

- Used only on the above mentioned crops and follow the recommended safety period before harvest.
- Do not allow animals to enter or graze in treated area.

**RE-ENTRY PERIOD:**

12 hours from last application.

**COMPATIBILITY:**

Compatible with most commonly used fungicides, insecticides, micronutrients and growth regulators, but compatibility test is recommended before application.

• For more details about first aid & precautions please refer to first aid & precautions index.